

# NONPARAMETRIC ESTIMATION OF MULTIVARIATE CONVEX-TRANSFORMED DENSITIES

BY ARSENI SEREGIN<sup>\*,†</sup> AND JON A. WELLNER<sup>†,‡</sup>

*University of Washington<sup>‡</sup>*

We study estimation of multivariate densities  $p$  of the form  $p(x) = h(g(x))$  for  $x \in \mathbb{R}^d$  and for a fixed function  $h$  and an unknown convex function  $g$ . The canonical example is  $h(y) = e^{-y}$  for  $y \in \mathbb{R}$ ; in this case the resulting class of densities

$$\mathcal{P}(e^{-y}) = \{p = \exp(-g) : g \text{ is convex}\}$$

is well-known as the class of *log-concave* densities. Other functions  $h$  allow for classes of densities with heavier tails than the log-concave class.

We first investigate when the MLE  $\hat{p}$  exists for the class  $\mathcal{P}(h)$  for various choices of monotone transformations  $h$  including decreasing and increasing functions  $h$ . The resulting models for increasing transformations  $h$  extend the classes of *log-convex* densities studied previously in the econometrics literature corresponding to  $h(y) = \exp(y)$ .

We then establish consistency of the MLE for fairly general functions  $h$ , including the log-concave class  $\mathcal{P}(e^{-y})$  and many others. In a final section we provide asymptotic minimax lower bounds for estimation of  $p$  and its vector of derivatives at a fixed point  $x_0$  under natural smoothness hypotheses on  $h$  and  $g$ . The proofs rely heavily on results from convex analysis.

## 1. Introduction and Background.

1.1. *Log-concave and  $r$ -concave densities.* A probability density  $p$  on  $\mathbb{R}^d$  is called log-concave if it can be written as

$$p(x) = \exp(-g(x))$$

for some convex function  $g : \mathbb{R}^d \rightarrow (-\infty, \infty]$ . We let  $\mathcal{P}(e^{-y})$  denote the class of all log-concave densities on  $\mathbb{R}^d$ . As shown by [Ibragimov \[1956\]](#), a

---

\*Research supported in part by NSF grant DMS-0503822

†Research supported in part by NSF grant DMS-0804587 and NI-AID grant 2R01 AI291968-04

AMS 2000 subject classifications: Primary 62G07, 62H12; secondary 62G05, 62G20

Keywords and phrases: consistency, log-concave density estimation, lower bounds, maximum likelihood, mode estimation, nonparametric estimation, qualitative assumptions, shape constraints, strongly unimodal, unimodal

density function  $p$  on  $\mathbb{R}$  is log-concave if and only if its convolution with any unimodal density is again unimodal.

Log-concave densities have proved to be useful in a wide range of statistical problems: see [Walther \[2010\]](#) for a survey of recent developments and statistical applications of log-concave densities on  $\mathbb{R}$  and  $\mathbb{R}^d$ , and see [Cule, Samworth and Stewart \[2007\]](#) for several interesting applications of estimators of such densities in  $\mathbb{R}^d$ .

Because the class of multivariate log-concave densities contains the class of multivariate normal densities and is preserved under a number of important operations (such as convolution and marginalization), it serves as a valuable nonparametric surrogate or replacement for the class of normal densities. Further study of the class of log-concave from this perspective has been given by [Schuhmacher, Hüsler and Duembgen \[2009\]](#). On the analysis side, various isoperimetric and Poincaré type inequalities have been obtained by [Bobkov \[1999, 2007b\]](#), [Fougères \[2005\]](#), and [Milman and Sodin \[2008\]](#).

Log-concave densities have the slight drawback that the tails must be decreasing exponentially, so a number of authors, including [Koenker and Mizera \[2008\]](#), have proposed using generalizations of the log-concave family involving  $r$ -concave densities defined as follows. For  $a, b \in \mathbb{R}$ ,  $r \in \mathbb{R}$ , and  $\lambda \in (0, 1)$ , define the generalized mean of order  $r$ ,  $M_r(a, b; \lambda)$ , by

$$M_r(a, b; \lambda) = \begin{cases} ((1 - \lambda)a^r + \lambda b^r)^{1/r}, & r \neq 0, \ a, b > 0, \\ 0, & r < 0, \ ab = 0, \\ a^{1-\lambda}b^\lambda, & r = 0. \end{cases}$$

Then a density function  $p$  is  $r$ -concave on  $C \subset \mathbb{R}^d$  if and only

$$p((1 - \lambda)x + \lambda y) \geq M_r(p(x), p(y); \lambda) \quad \text{for all } x, y \in C, \ \lambda \in (0, 1).$$

We denote the class of all  $r$ -concave densities on  $C \subset \mathbb{R}^d$  by  $\widehat{\mathcal{P}}(y_+^{1/r}; C)$ , and write  $\widehat{\mathcal{P}}(y_+^{1/r})$  when  $C = \mathbb{R}^d$ . As noted by [Dharmadhikari and Joag-Dev \[1988\]](#), page 86, for  $r \leq 0$  it suffices to consider  $\widehat{\mathcal{P}}(y_+^{1/r})$ , and it is almost immediate from the definitions that  $p \in \widehat{\mathcal{P}}(y_+^{1/r})$  if and only if  $p(x) = (g(x))^{1/r}$  for some convex function  $g$  from  $\mathbb{R}^d$  to  $[0, \infty)$ . For  $r > 0$ ,  $p \in \widehat{\mathcal{P}}(y_+^{1/r}; C)$  if and only if  $p(x) = (g(x))^{1/r}$  where  $g$  mapping  $C$  into  $(0, \infty)$  is concave.

These results motivate definitions of the classes  $\mathcal{P}(y_+^{-s}) = \{p(x) = g(x)^{-s} : g \text{ is convex}\}$  for  $s \geq 0$  and, more generally, for a fixed monotone function  $h$  from  $\mathbb{R}$  to  $\mathbb{R}$ ,

$$\mathcal{P}(h) \equiv \{p(x) = h(g(x)) : g \text{ convex}\}.$$

Such generalizations of log-concave densities and log-concave measures based on means of order  $r$  have been introduced by a series of authors, sometimes with differing terminology, apparently starting with [Avriel \[1972\]](#), and continuing with [Borell \[1975\]](#), [Brascamp and Lieb \[1976\]](#), [Prékopa \[1973\]](#), [Rinott \[1976\]](#), and [Uhrin \[1984\]](#). A nice summary of these connections is given by [Dharmadhikari and Joag-Dev \[1988\]](#). These authors also present results concerning preservation of  $r$ -concavity under a variety of operations including products, convolutions, and marginalization. In the mathematics literature, the underlying fundamental inequality has come to be known as the Borell-Brascamp-Lieb inequality; see e.g. [Cordero-Erausquin, McCann and Schmuckenschläger \[2001\]](#). For these heavy-tailed classes, development of isoperimetric and Poincaré inequalities is also underway: see e.g. [Bobkov \[2007a\]](#) and [Bobkov and Ledoux \[2009\]](#).

Despite the long-standing and current rapid development of the properties of such classes of densities on the probability side, very little has been done from the standpoint of nonparametric estimation, especially when  $d \geq 2$ .

Nonparametric estimation of a log-concave density on  $\mathbb{R}^d$  was initiated by [Cule, Samworth and Stewart \[2007\]](#). These authors developed an algorithm for computing their estimators and explored several interesting applications. [Koenker and Mizera \[2008\]](#) developed a family of penalized criterion functions related to the Rényi divergence measures, and explored duality in the optimization problems. They did not succeed in establishing consistency of their estimators, but did investigate Fisher consistency. Recently, [Cule and Samworth \[2009\]](#) have established consistency of the (nonparametric) maximum likelihood estimator of a log-concave density on  $\mathbb{R}^d$ , even in a setting of model misspecification: when the true density is not log-concave, then the estimator converges to the closest log-concave density to the true density in the sense of Kullback-Leibler divergence.

In this paper our goal is to investigate maximum likelihood estimation in the classes  $\mathcal{P}(h)$  corresponding to a fixed monotone (decreasing or increasing) function  $h$ . In particular, for decreasing functions  $h$ , we handle all the  $r$ -concave classes  $\mathcal{P}(y_+^{1/r})$  with  $r = -1/s$  and  $r \leq -1/d$  (or  $s \geq d$ ). On the increasing side, we treat, in particular, the cases  $h(y) = y1_{[0,\infty)}(y)$  and  $h(y) = e^y$  with  $C = \mathbb{R}_+^d$ . The first of these corresponds to an interesting class of models which can be thought of as multivariate generalizations of the class of decreasing and convex densities on  $\mathbb{R}_+$  treated by [Groeneboom, Jongbloed and Wellner \[2001\]](#), while the second,  $h(y) = e^y$  corresponds to multivariate versions of the log-convex families studied by [An \[1998\]](#). Note that our increasing classes  $\mathcal{P}(y_+^{1/r}, \mathbb{R}_+^d)$  with  $r > 0$  are quite different from the  $r$ -concave classes defined above and appear to be completely new, cor-

responding instead to  $r$ -convex densities on  $\mathbb{R}_+^d$ .

Here is an outline of the rest of the paper. All of our main results are presented in Section 2. Subsection 2.1 gives definitions and basic properties of the transformations involved. Subsection 2.2 establishes existence of the maximum likelihood estimators for both increasing and decreasing transformations  $h$  under suitable conditions on the function  $h$ . In subsection 2.3 we give statements concerning consistency of the estimators, both in the Hellinger metric and in uniform metrics under natural conditions. In subsection 2.4 we present asymptotic minimax lower bounds for estimation in these classes under natural curvature hypotheses. We conclude this section with a brief discussion of conjectures concerning attainability of the minimax rates by the maximum likelihood estimators. All the proofs are given in Section 3. We summarize a number of key results from convex analysis in an appendix, Section 4.

**1.2. Convex-transformed density estimation.** Now let  $X_1, \dots, X_n$  be  $n$  independent random variables distributed according to a probability density  $p_0 = h(g_0(x))$  on  $\mathbb{R}^d$  here  $h$  is a fixed monotone (increasing or decreasing) function and  $g_0$  is an (unknown) convex function. The probability measure on the Borel sets  $\mathcal{B}_d$  corresponding to  $p_0$  is denoted by  $P_0$ .

The maximum likelihood estimator (MLE) of a log-concave density on  $\mathbb{R}$  was introduced in Rufibach [2006] and Dümbgen and Rufibach [2007]. Algorithmic aspects were treated in Rufibach [2007] and in a more general framework in Dümbgen, Hüsler and Rufibach [2007], while consistency with respect to the Hellinger metric was established by Pal, Woodroffe and Meyer [2007], and rates of convergence of  $\hat{f}_n$  and  $\hat{F}_n$  were established by Dümbgen and Rufibach [2007]. Asymptotic distribution theory for the MLE of a log-concave density on  $\mathbb{R}$  was established by Balabdaoui, Rufibach and Wellner [2009].

If  $\mathcal{C}$  denotes the class of all closed proper convex functions  $g : \mathbb{R}^d \rightarrow (-\infty, \infty]$ , the estimator  $\hat{g}_n$  of  $g_0$  is the maximizer of the functional:

$$\mathbb{L}_n g \equiv \int (\log h) \circ g d\mathbb{P}_n$$

over the class  $\mathcal{G}(h) \subset \mathcal{C}$  of all convex functions  $g$  such that  $h \circ g$  is a density and where  $\mathbb{P}_n$  is the empirical measure of the observations. The maximum likelihood estimator of the convex-transformed density  $p_0$  is then  $\hat{p}_n := h(\hat{g}_n)$  when it exists and is unique. We investigate conditions for existence and uniqueness in Section 2.

## 2. Main results.

**2.1. Definitions and basic properties.** To construct the classes of convex-transformed densities of interest here, we first need to define two classes of monotone transformations. An *increasing transformation*  $h$  is a nondecreasing function  $\mathbb{R} \rightarrow \mathbb{R}_+$  such that  $h(-\infty) = 0$  and  $h(+\infty) = +\infty$ . We define the limit points  $y_0 < y_\infty$  of the increasing transformation  $h$  by:

$$y_0 = \inf\{y : h(y) > 0\}$$

$$y_\infty = \sup\{y : h(y) < +\infty\}.$$

We make the following assumptions about the asymptotic behavior of the increasing transformation.

- (I.1) The function  $h(y)$  is  $o(|y|^{-\alpha})$  for some  $\alpha > d$  as  $y \rightarrow -\infty$ .
- (I.2) If  $y_\infty < +\infty$  then  $h(y) \asymp (y_\infty - y)^{-\beta}$  for some  $\beta > d$  as  $y \uparrow y_\infty$ .
- (I.3) The function  $h$  is continuously differentiable on the interval  $(y_0, y_\infty)$ .

Note that the assumption I.1 is satisfied if  $y_0 > -\infty$ .

**DEFINITION 2.1.** For an increasing transformation  $h$  an increasing class of convex-transformed densities or simply an increasing model  $\mathcal{P}(h)$  on  $\mathbb{R}_+^d$  is the family of all bounded densities which have the form  $h \circ g \equiv h(g(\cdot))$ , where  $g$  is a closed proper convex function with  $\text{dom } g = \mathbb{R}_+^d$ .

**REMARK 2.2.** Consider a density  $h \circ g$  from an increasing model  $\mathcal{P}(h)$ . Since  $h \circ g$  is bounded we have  $g < y_\infty$ . The function  $\tilde{g} = \max(g, y_0)$  is convex and  $h \circ \tilde{g} = h \circ g$ . Thus we can assume that  $g \geq y_0$ .

A *decreasing transformation*  $h$  is a nonincreasing function  $\mathbb{R} \rightarrow \mathbb{R}_+$  such that  $h(-\infty) = +\infty$  and  $h(+\infty) = 0$ . We define the limit points  $y_0 > y_\infty$  of the decreasing transformation  $h$  by:

$$y_0 = \sup\{y : h(y) > 0\}$$

$$y_\infty = \inf\{y : h(y) < +\infty\}.$$

We make the following assumptions about the asymptotic behavior of the decreasing transformation.

- (D.1) The function  $h(y)$  is  $o(y^{-\alpha})$  for some  $\alpha > d$  as  $y \rightarrow +\infty$ .
- (D.2) If  $y_\infty > -\infty$  then  $h(y) \asymp (y - y_\infty)^{-\beta}$  for some  $\beta > d$  as  $y \downarrow y_\infty$ .
- (D.3) If  $y_\infty = -\infty$  then  $h(y)^\gamma h(-Cy) = o(1)$  for some  $\gamma, C > 0$  as  $y \rightarrow -\infty$ .

(D.4) The function  $h$  is continuously differentiable on the interval  $(y_\infty, y_0)$ .

Note that the assumption D.1 is satisfied if  $y_0 < +\infty$ . Now we define the decreasing class of densities  $\mathcal{P}(h)$ .

DEFINITION 2.3. *For a decreasing transformation  $h$  a decreasing class of convex-transformed densities or simply a decreasing model  $\mathcal{P}(h)$  on  $\mathbb{R}^d$  is the family of all bounded densities which have the form  $h \circ g$ , where  $g$  is a closed proper convex function with  $\dim(\text{dom } g) = d$ .*

REMARK 2.4. *Consider a density  $h \circ g$  from a decreasing model  $\mathcal{P}(h)$ . Since  $h \circ g$  is bounded we have  $g > y_\infty$ . For the sublevel set  $C = \text{lev}_{y_0} g$ , the function  $\tilde{g} = g + \delta(\cdot | C)$  is convex and  $h \circ \tilde{g} = h \circ g$ . Thus we can assume that  $\text{lev}_{y_0} h = \text{dom } g$ .*

For a monotone transformation  $h$  we denote by  $\mathcal{G}(h)$  the class of all convex functions  $g$  such that  $h \circ g$  belongs to a decreasing class  $\mathcal{P}(h)$ . The following lemma allows us to compare models defined by increasing or decreasing transformations  $h$

LEMMA 2.5. *Consider two decreasing (or increasing) models  $\mathcal{P}(h_1)$  and  $\mathcal{P}(h_2)$ . If  $h_1 = h_2 \circ f$  for some convex function  $f$  then  $\mathcal{P}(h_1) \subseteq \mathcal{P}(h_2)$ .*

PROOF. The argument below is for a decreasing model. For an increasing model the proof is similar. If  $f(x) > f(y)$  for some  $x < y$  then  $f$  is decreasing on  $(-\infty, x)$ ,  $f(-\infty) = +\infty$ , and therefore  $h_2$  is constant on  $(f(x), +\infty)$  and we can redefine  $f(y) = f(x)$  for all  $y < x$ . Thus we can always assume that  $f$  is nondecreasing.

For any convex function  $g$ , the function  $f \circ g$  is also convex. Therefore, if  $p = h_1 \circ g \in \mathcal{P}(h_1)$ , then  $p = h_2 \circ f \circ g \in \mathcal{P}(h_2)$ .  $\square$

In this section we discuss several examples of monotone models. First two families based on increasing transformations  $h$ :

EXAMPLE 2.6 (Log-convex densities). *This increasing model is defined by  $h(y) = e^y$  and  $C = \mathbb{R}_+^d$ . Limit points are  $y_0 = -\infty$  and  $y_\infty = \infty$ . Assumption I.1 holds for any  $\alpha > d$ . These classes of densities were considered by An [1998], who established several useful preservation properties. In particular, log-convexity is preserved under mixtures (An [1998, Proposition 3]) and under marginalization (An [1998, Remark 8, p. 361]).*

EXAMPLE 2.7 ( $r$ -Convex densities). *This family of increasing models is defined by the transforms  $h(y) = \max(y, 0)^s = y_+^s$  with  $s > 0$  and  $C = \mathbb{R}_+^d$ . Limit points are:  $y_0 = 0$  and  $y_\infty = +\infty$ . Assumption I.1 holds for any  $\alpha > d$ . As noted in Section 1 the model  $\mathcal{P}(y_+^{1/r}, \mathbb{R}_+^d)$  corresponds to the class of  $r$ -convex densities, with  $r = \infty$  corresponding to the log-convex densities of the previous example. For  $r < \infty$  these classes seem to have not been previously discussed or considered, except in special cases: the case  $r = 1$  and  $d = 1$  corresponds to the class of decreasing convex densities on  $\mathbb{R}_+$  considered by [Groeneboom, Jongbloed and Wellner \[2001\]](#). It follows from Lemma 2.5 that*

$$(2.1) \quad \mathcal{P}(e^y, \mathbb{R}_+^d) \subset \mathcal{P}(y_+^{s_2}, \mathbb{R}_+^d) \subset \mathcal{P}(y_+^{s_1}, \mathbb{R}_+^d), \quad \text{for } 0 < s_2 < s_1 < \infty.$$

Now for some models based on decreasing transformations  $h$ :

EXAMPLE 2.8 (Log-concave densities). *This decreasing model is defined by the transform  $h(y) = e^{-y}$ . Limit points are:  $y_0 = +\infty$  and  $y_\infty = -\infty$ . Assumption D.1 holds for any  $\alpha > d$ . Assumption D.3 holds for any  $\gamma > C > 0$ .*

Many parametric models are subsets of this model. Below we specify convex functions  $g$  which correspond to the densities of several distributions:

1. *Uniform:* Density of a uniform distribution on a convex set  $C$  is log-concave:

$$g(x) = -\log(\mu[C]) + \delta(x | C).$$

2. *Normal:* Density of a multivariate normal distribution  $(\mu, \Sigma)$  with  $\Sigma$  nonsingular is log-concave:

$$g(x) = \frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu) + \frac{1}{2} \log |\Sigma| + \frac{d}{2} \log 2\pi.$$

3. *Gamma:* Density of Gamma distribution  $(r, \lambda)$  is log-concave for  $r > 1$ :

$$g(x) = -(r - 1) \log x + \lambda x - r \log \lambda + \log \Gamma(r).$$

4. *Beta:* Density of Beta distribution  $(\alpha, \beta)$  is log-concave for  $\alpha, \beta > 1$ :

$$g(x) = -(\alpha - 1) \log x - (\beta - 1) \log(1 - x) + \log B(\alpha, \beta).$$

Gumbel, Fréchet and logistic distributions also have log-concave densities.

EXAMPLE 2.9 ( $r$ -Concave densities; Power-convex densities). *This family of decreasing models is defined by the transforms  $h(y) = y_+^{-s}$  for  $s > d$ . Limit points are:  $y_0 = +\infty$  and  $y_\infty = 0$ . Assumption D.1 holds for any  $\alpha \in (d, s]$ . Assumption D.2 holds for  $\beta = s$ . As noted in Section 1 the model  $\mathcal{P}(y_+^{1/r}) = \mathcal{P}(y_+^{-s})$  (with  $r = -1/s < 0$ ) corresponds to the class of  $r$ -concave densities. From Lemma 2.5 we have the following inclusion:*

$$(2.2) \quad \mathcal{P}(e^{-y}) \subset \mathcal{P}(y_+^{-s_2}) \subset \mathcal{P}(y_+^{-s_1}), \quad \text{for } s_1 < s_2.$$

The models defined by power transformations include some parametric models with heavier than exponential tails. The following examples are discussed in Borell [1975]. We use Johnson and Kotz [1972] as a reference.

1. *Pareto* (Johnson and Kotz [1972] 42.3): Density of a multivariate Pareto distribution  $(\theta, a)$  is power-convex for  $s \in (d, a + d]$ :

$$g(x) = \left( \frac{\Gamma(a) \prod \theta_i}{\Gamma(a + d)} \right)^{1/s} \left( 1 - d + \sum x_i / \theta_i \right)^{(a+d)/s}.$$

2. *Student* (Johnson and Kotz [1972] 37.3): Density of a multivariate  $t$ -distribution  $(d, n, \mu, \Sigma)$  is power-convex for  $s \in (d, n + d]$ :

$$g(x) = \left( \frac{\Gamma(n/2)(n\pi)^{d/2} |\Sigma|^{1/2}}{\Gamma((d+n)/2)} \right)^{1/s} \left( 1 + \frac{1}{n} (x - \mu)^T \Sigma^{-1} (x - \mu) \right)^{(d+n)/2s}.$$

3. *Snedecor* (Johnson and Kotz [1972] 40.8): Density of a multivariate  $F$ -distribution  $(n_0, n_1, \dots, n_d)$  with  $n_i \geq 2$  is power-convex for  $s \in (d, (n_0/2) + d]$ :

$$g(x) = C(n_i, d, s) \left( n_0 + \sum_{i=1}^d n_i x_i \right)^{n/2s} \left( \prod_{i=1}^d x_i^{2-n_i} \right)^{1/2s},$$

where  $n = \sum_{i=0}^d n_i$ .

Since the distributions above belong to the power-convex models only for bounded values of the parameter  $s$  the inclusion (2.2) implies that they do not belong to the log-concave model (corresponding to  $s = +\infty$ )

Borell [1975] developed a framework which unifies log-concave and power-convex densities and gives interesting characterization for these classes. Here we briefly state the main result.



DEFINITION 2.10. Let  $C \subseteq \mathbb{R}^d$  be an open convex set and let  $s \in \overline{\mathbb{R}}$ . Then we define  $\mathcal{M}_s(C)$  as the family of all positive Radon measures  $\mu$  on  $C$  such that:

$$(2.3) \quad \mu_*(\theta A + (1 - \theta)B) \geq [\theta \mu_*(A)^s + (1 - \theta) \mu_*(B)^s]^{1/s}$$

holds for all  $\emptyset \neq A, B \subseteq C$  and all  $\theta \in (0, 1)$ . We define  $\mathcal{M}_s^\circ(C)$  as a subfamily of  $\mathcal{M}_s(C)$  which consists of probability measures such that the affine hull of its support has dimension  $d$ . Here  $\mu_*$  is the inner measure corresponding to  $\mu$  and the cases  $s = 0, \infty$  are defined by continuity.

Then one of the main results of Borell [1975], Prékopa [1973], and Rinott [1976] is as follows:

THEOREM 2.11 (Borell, Prekopa, Rinott). For  $s < 0$  the family  $\mathcal{M}_s^\circ(\mathbb{R}^d)$  coincides with the power-convex family  $\mathcal{P}(y_+^{-d+1/s})$ . For  $s = 0$  the family  $\mathcal{M}_0^\circ(\mathbb{R}^d)$  coincides with the log-concave family  $\mathcal{P}(e^{-y})$ .

Finally, Corollary 3.1 Borell [1975] says that the condition (2.3) can be relaxed:

THEOREM 2.12 (Borell). Let  $\Omega \subseteq \mathbb{R}^d$  be an open convex set and let  $\mu$  be a positive Radon measure in  $\Omega$ . Then  $\mu \in \mathcal{M}_s(\Omega)$  if and only if

$$\mu\left(\frac{1}{2}A_1 + \frac{1}{2}A_2\right) \geq \left(\frac{1}{2}\mu(A_1)^s + \frac{1}{2}\mu(A_2)^s\right)^{1/s}$$

holds for all compact (or open, or semiopen) blocks  $A_1, A_2 \subseteq \Omega$  (i.e. rectangles with sides parallel to the coordinate axes).

Theorem 2.11 gives a special case of what has come to be known as the Borell-Brascamp-Lieb inequality; see e.g. Dharmadhikari and Joag-Dev [1988], and Brascamp and Lieb [1976]. The current terminology is apparently due to Cordero-Erausquin, McCann and Schmuckenschläger [2001].

2.2. *Existence of the Maximum Likelihood Estimators.* Now suppose that  $X_1, \dots, X_n$  are i.i.d. with density  $p_0(x) = h(g_0(x))$  for a fixed monotone transformation  $h$  and a convex function  $g_0$ . As before,  $\mathbb{P}_n = n^{-1} \sum_{i=1}^n \delta_{X_i}$  is the empirical measure of the  $X_i$ 's and  $P_0$  is the probability measure corresponding to  $p_0$ . Then  $\mathbb{L}_n g = \mathbb{P}_n \log h \circ g$  is the log-likelihood function (divided by  $n$ ), and

$$\hat{p}_n \equiv \operatorname{argmax}\{\mathbb{L}_n g : h \circ g \in \mathcal{P}(h)\}$$

is the *maximum likelihood estimator* of  $p$  over the class  $\mathcal{P}(h)$ , assuming it exists and is unique. We also write  $\hat{g}_n$  for the MLE of  $g$ . We first state our main results concerning existence and uniqueness of the MLE's for the classes  $\mathcal{P}(h)$ :

**THEOREM 2.13.** *Suppose that  $h$  is an increasing transformation satisfying assumptions [I.1](#) - [I.3](#). Then the MLE  $\hat{p}_n$  exists almost surely for the model  $\mathcal{P}(h)$ .*

**THEOREM 2.14.** *Suppose that  $h$  is a decreasing transformation satisfying assumptions [D.1](#) - [D.4](#). Then the MLE  $\hat{p}_n$  exists almost surely for the model  $\mathcal{P}(h)$  if*

$$n \geq n_d \equiv \begin{cases} d(1 + \gamma), & \text{if } y_\infty = -\infty \\ d + \frac{\beta d^2}{\alpha(\beta - d)}, & \text{if } y_\infty > -\infty. \end{cases}$$

Uniqueness of the MLE is known for the log-concave model  $\mathcal{P}(e^{-y})$ , see e.g. [Duembgen and Rufibach \[2009\]](#) for  $d = 1$  and [Cule et al. \[2007\]](#) for  $d \geq 1$ . For a brief further comment see section [2.5](#).

**2.3. Consistency of the Maximum Likelihood Estimators.** Once existence of the MLE's is assured, our attention shifts to other properties of the estimators: our main concern in this sub-section is consistency. While for a decreasing model it is possible to prove consistency without any restrictions, for an increasing model we need the following assumptions about the true density  $h \circ g_0$ :

**(I.4)** The function  $g_0$  is bounded by some constant  $C < y_\infty$ .

**(I.5)** If  $d > 1$  then we have, with  $|x| \equiv \prod_{j=1}^d x_j$  for  $x \in \mathbb{R}_+^d$ ,

$$C_g \equiv \int_{\mathbb{R}_+^d} \log \left( \frac{1}{|x| \wedge 1} \right) dP_0(x) < \infty.$$

**REMARK 2.15.** *Note that for  $d = 1$  the assumption [I.5](#) follows from the assumption [I.4](#) and integrability of  $\log(1/x)$  at zero. This assumption is also true if  $P$  has finite marginal densities.*

**(I.6)** We have:

$$\int_{\mathbb{R}_+^d} (h \log h) \circ g_0(x) dx < \infty.$$

Let  $H(p, q)$  denote the Hellinger distance between two probability measures with densities  $p$  and  $q$  with respect to Lebesgue measure on  $\mathbb{R}^d$ :

$$(2.4) \quad H^2(p, q) \equiv \frac{1}{2} \int_{\mathbb{R}^d} (\sqrt{p}(x) - \sqrt{q}(x))^2 dx = 1 - \int_{\mathbb{R}^d} \sqrt{p(x)q(x)} dx.$$

Our main results about increasing models are as follows.

**THEOREM 2.16.** *For an increasing model  $\mathcal{P}(h)$  where  $h$  satisfies assumptions [I.1](#) - [I.3](#) and for the true density  $h \circ g_0$  which satisfies assumptions [I.4](#) - [I.6](#), the sequence of MLEs  $\{\hat{p}_n = h \circ \hat{g}_n\}$  is Hellinger consistent:*

$$H(\hat{p}_n, p_0) = H(h \circ \hat{g}_n, h \circ g_0) \rightarrow_{a.s.} 0.$$

**THEOREM 2.17.** *For an increasing model  $\mathcal{P}(h)$  where  $h$  satisfies assumptions [I.1](#) - [I.3](#) and for the true density  $h \circ g_0$  which satisfies assumptions [I.4](#) - [I.6](#), the sequence of MLEs  $\hat{g}_n$  is pointwise consistent. That is  $\hat{g}_n(x) \rightarrow_{a.s.} g_0(x)$  for  $x \in \text{ri}(\mathbb{R}_+^d)$  and convergence is uniform on compacta.*

The results about decreasing models can be formulated in a similar way.

**THEOREM 2.18.** *For a decreasing model  $\mathcal{P}(h)$  where  $h$  satisfies assumptions [D.1](#) - [D.4](#), the sequence of MLEs  $\{\hat{p}_n = h \circ \hat{g}_n\}$  is Hellinger consistent:*

$$H(\hat{p}_n, p_0) = H(h \circ \hat{g}_n, h \circ g_0) \rightarrow_{a.s.} 0.$$

**THEOREM 2.19.** *For a decreasing model  $\mathcal{P}(h)$  with  $h$  satisfying assumptions [D.1](#) - [D.4](#), the sequence of MLEs  $\hat{g}_n$  is pointwise consistent in the following sense. Define  $g_0^* = g_0 + \delta(\cdot | \text{ri}(\text{dom } g_0))$ . Then  $g_0^* = g_0$  a.e.,*

$$\hat{g}_n \rightarrow_{a.s.} g_0^*,$$

*and the convergence is uniform on compacta. Moreover, if  $\text{dom } g_0 = \mathbb{R}^d$  then:*

$$\|h \circ \hat{g}_n - h \circ g_0\|_\infty \rightarrow_{a.s.} 0.$$

**2.4. Local Asymptotic Minimax Lower Bounds.** In this section establish local asymptotic minimax lowers for any estimator of several functionals of interest on the family  $\mathcal{P}(h)$  of convex transformed densities. We start with several general results following [Jongbloed \[2000\]](#), and then apply them to estimation at a point fixed point and to mode estimation.

First, we define minimax risk as in [Donoho and Liu \[1991\]](#):

DEFINITION 2.20. Let  $\mathcal{P}$  be a class of densities on  $\mathbb{R}^d$  with respect to Lebesgue measure and  $T$  be a functional  $T : \mathcal{P} \rightarrow \mathbb{R}$ . For an increasing convex loss function  $l$  on  $\mathbb{R}_+$  we define the minimax risk as:

$$(2.5) \quad R_l(n; T, \mathcal{P}) = \inf_{t_n} \sup_{p \in \mathcal{P}} \mathbb{E}_{p^{\times n}} l(|t_n(X_1, \dots, X_n) - Tp|),$$

where  $t_n$  ranges over all possible estimators of  $Tp$  based on  $X_1, \dots, X_n$ .

The main result (Theorem 1) in Jongbloed [2000] can be formulated as:

THEOREM 2.21 (Jongbloed). Let  $\{p_n\}$  be a sequence of densities in  $\mathcal{P}$  such that:

$$\limsup_{n \rightarrow \infty} \sqrt{n} H(p_n, p) \leq \tau$$

for some density  $p$  in  $\mathcal{P}$ . Then:

$$(2.6) \quad \liminf_{n \rightarrow \infty} \frac{R_l(n; T, \{p, p_n\})}{l((1/4)e^{-2\tau^2}|T(p_n) - T(p)|)} \geq 1.$$

It will be convenient to reformulate this result in the following form:

COROLLARY 2.22. Suppose that for any  $\varepsilon > 0$  small enough, there exists  $p_\varepsilon \in \mathcal{P}$  such that for some  $r > 0$ :

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} |Tp_\varepsilon - Tp| = 1$$

and

$$\limsup_{\varepsilon \rightarrow 0} \varepsilon^{-r} H(p_\varepsilon, p) \leq c.$$

Then, there exists a sequence  $\{p_n\}$  such that:

$$(2.7) \quad \liminf_{n \rightarrow \infty} n^{1/2r} R_1(n; T, \{p, p_n\}) \geq \frac{1}{4(2re)^{1/2r}} c^{-1/r},$$

where  $R_1$  is the risk which corresponds to  $l(x) = |x|$ .

Corollary 2.22 shows that for a fixed change in the value of the functional  $T$ , a family  $p_\varepsilon$  which is closer to the true density  $p$  with respect to Hellinger distance provides sharper lower bound. This suggests that for the functional  $T$  which depends only on the local structure of the density we would like our family  $\{p_\varepsilon\}$  to deviate from  $p$  also locally. Below, we define formally such local deviations.

DEFINITION 2.23. *We call a family of measurable functions  $\{p_\varepsilon\}$  a deformation of a measurable function  $p$  if  $p_\varepsilon$  is defined for any  $\varepsilon > 0$  small enough,*

$$\lim_{\varepsilon \rightarrow 0} \text{ess sup } |p - p_\varepsilon| = 0,$$

*and there exists a bounded family of real numbers  $r_\varepsilon$  and a point  $x_0$  such that:*

$$\begin{aligned} \mu[\text{supp } |p_\varepsilon(x) - p(x)|] &> 0, \\ \text{supp } |p_\varepsilon(x) - p(x)| &\subseteq B(x_0, r_\varepsilon). \end{aligned}$$

*If in addition we have:*

$$\lim_{\varepsilon \rightarrow 0} r_\varepsilon = 0$$

*we say that  $\{p_\varepsilon\}$  is a local deformation at  $x_0$ .*

Since for a deformation  $p_\varepsilon$  we have  $\mu[\text{supp } |p_\varepsilon(x) - p(x)|] > 0$ , for every  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $\mu\{x : |p_\varepsilon(x) - p(x)| > \delta\} > 0$  and thus the  $L_r$ -distance from  $p_\varepsilon$  to  $p$  is positive for all  $\varepsilon > 0$ . Note that this is always true if  $p$  and  $p_\varepsilon$  are continuous at  $x_0$  and  $p_\varepsilon(x_0) \neq p(x_0)$ .

Now we can state our lower bound for estimation of the convex-transformed density value at a fixed point  $x_0$ . This result relies on the properties of strongly convex functions as described in Appendix 4.4 and can be applied to both increasing and decreasing classes of convex-transformed densities.

THEOREM 2.24. *Let  $h$  be a monotone transformation, let  $p = h \circ g \in \mathcal{P}(h)$  be a convex-transformed density, and suppose that  $x_0$  is a point in  $\text{ri}(\text{dom } g)$  such that  $h$  is continuously differentiable at  $g(x_0)$ ,  $h \circ g(x_0) > 0$ ,  $h' \circ g(x_0) \neq 0$ , and  $\text{curv}_{x_0} g > 0$ . Then, for the functional  $T(h \circ g) \equiv g(x_0)$  there exists a sequence  $\{p_n\} \subset \mathcal{P}(h)$  such that:*

$$(2.8) \quad \liminf_{n \rightarrow \infty} n^{\frac{2}{d+4}} R_1(n; T, \{h \circ g, p_n\}) \geq C(d) \left[ \frac{h \circ g(x_0)^2 \text{curv}_{x_0} g}{h' \circ g(x_0)^4} \right]^{\frac{1}{d+4}},$$

*where the constant  $C(d)$  depends only on the dimension  $d$ .*

REMARK 2.25. *If in addition  $g$  is twice continuously differentiable at  $x_0$  and  $\nabla^2 g(x_0)$  is positive definite, then by Lemma 4.22 we have  $\text{curv}_{x_0} g = \det(\nabla^2 g(x_0))$ .*

In [Jongbloed \[2000\]](#) lower bounds were constructed for functionals with values in  $\mathbb{R}$ . However, it is easy to see that the proof does not change for functionals with values in an arbitrary metric space  $(V, s)$  if instead of  $|Tp - Tp_n|$  we consider  $s(Tp, Tp_n)$ . We define:

$$(2.9) \quad R_s(n; T, \mathcal{P}) = \inf_{t_n} \sup_{p \in \mathcal{P}} E_{p^{\times n}} s(t_n(X_1, \dots, X_n), Tp),$$

and the analogue of Corollary [2.22](#) has the following form:

**COROLLARY 2.26.** *Suppose that for any  $\varepsilon > 0$  small enough, there exists  $p_\varepsilon \in \mathcal{P}$  such that for some  $r > 0$ :*

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} s(Tp_\varepsilon, Tp) &= 1 \\ \limsup_{\varepsilon \rightarrow 0} \varepsilon^{-r} H(p_\varepsilon, p) &\leq c. \end{aligned}$$

*Then, there exists a sequence  $\{p_n\}$  such that:*

$$(2.10) \quad \liminf_{n \rightarrow \infty} n^{1/2r} R_s(n; T, \{p, p_n\}) \geq \frac{1}{4(re)^{1/2r}} c^{-1/r}.$$

In this section we consider estimation of the functional

$$T(h \circ g) = \operatorname{argmin}(g) \in \mathbb{R}^d$$

for the density  $p = h \circ g \in \mathcal{P}(h)$  assuming that the minimum is unique. This is equivalent to estimation of the mode of  $p = h \circ g$ .

Construction of a lower bound for the functional  $T$  is similar to the procedure we presented for estimation of  $p = h \circ g$  at a fixed point  $x_0$ . Again, we use two opposite deformations: one is local and changes the functional value, another is a convex combination with a fixed deformation and negligible in Hellinger distance computation. However, in this case the minimax rate depends also on the growth rate of  $g$ .

**THEOREM 2.27.** *Let  $h$  be a decreasing transformation,  $h \circ g \in \mathcal{P}(h)$  be a convex-transformed density and a point  $x_0 \in \operatorname{ri}(\operatorname{dom} g)$  be a unique global minimum of  $g$  such that  $h$  is continuously differentiable at  $g(x_0)$ ,  $h' \circ g(x_0) \neq 0$  and  $\operatorname{curv}_{x_0} g > 0$ . In addition let us assume that  $g$  is locally Hölder continuous at  $x_0$ :*

$$|g(x) - g(x_0)| \leq L \|x - x_0\|^\gamma$$

with respect to some norm  $\|\cdot\|$ . Then, for the functional  $T(h \circ g) \equiv \operatorname{argmin} g$  there exists a sequence  $\{p_n\} \in \mathcal{P}(h)$  such that:

(2.11)

$$\liminf_{n \rightarrow \infty} n^{\frac{2}{\gamma(d+4)}} R_s(n; T, \{p, p_n\}) \geq C(d) L^{-\frac{1}{\gamma}} \left[ \frac{h \circ g(x_0)^2 \operatorname{curv}_{x_0} g}{h' \circ g(x_0)^4} \right]^{\frac{1}{\gamma(d+4)}},$$

where the constant  $C(d)$  depends only on the dimension  $d$  and metric  $s(x, y)$  is defined as  $\|x - y\|$ .

REMARK 2.28. If in addition  $g$  is twice continuously differentiable at  $x_0$  and  $\nabla^2 g(x_0)$  is positive definite, then by Lemma 4.22 we have  $\operatorname{curv}_{x_0} h = \det(\nabla^2 g(x_0))$  and  $g$  is locally Hölder continuous at  $x_0$  with exponent  $\gamma = 2$  and any constant  $L > \|\nabla^2 g(x_0)\|$ .

REMARK 2.29. Since  $\operatorname{curv}_{x_0} g > 0$  there exists constant  $C$  such that:

$$C\|x - x_0\|^2 \leq |g(x) - g(x_0)|,$$

and thus we have  $\gamma \in (0, 2]$ .

2.5. *Conjectures concerning uniqueness of MLEs.* There exist counterexamples to uniqueness for nonconvex transformations  $h$  which satisfy assumptions D.1 - D.4. They suggest that uniqueness of the MLE does not depend on the tail behavior of the transformation  $h$  but rather on the local properties of  $h$  in neighborhoods of the optimal values  $\hat{g}_n(X_i)$ . We conjecture that uniqueness holds for all monotone models if  $h$  is convex and  $h/|h'|$  is nondecreasing convex. Further work on the uniqueness issues is needed.

2.6. *Conjectures about rates of convergence for the MLEs.* We conjecture that the (optimal) rate of convergence  $n^{2/(d+4)}$  appearing in Theorem 2.24 for estimation of  $f(x_0)$  will be achieved by the MLE only for  $d = 2, 3$ . For  $d = 4$ , we conjecture that the MLE will come within a factor  $(\log n)^{-\gamma}$  (for some  $\gamma > 0$ ) of achieving the rate  $n^{1/4}$ , but for  $d > 4$  we conjecture that the rate of convergence will be the suboptimal rate  $n^{1/d}$ . This conjectured rate sub-optimality raises several interesting further issues:

- Can we find alternative estimators (perhaps via penalization or sieve methods) which achieve the optimal rates of convergence?
- For interesting sub-classes do maximum likelihood estimators remain rate optimal?

Baraud and Birgé [2009] have recently studied estimation in generalizations of our classes  $\mathcal{P}(h)$  in which both  $h$  and  $g$  are unknown with  $h : \mathbb{R}^m \rightarrow \mathbb{R}$  and  $g : \mathbb{R}^d \rightarrow \mathbb{R}^m$ . While they achieve optimal rates of estimation in their more general problem with a class of estimators based on multiple testing and model selection, their procedures will very likely have several drawbacks relative to the methods we have studied here, including (a) not belonging to the the classes of densities defined by the models, and (b) difficulties in computation or implementation.

### 3. Proofs.

#### 3.1. Preliminaries: properties of increasing transformations.

LEMMA 3.1. *Let  $h$  be a increasing transformation and  $g$  be a closed proper convex function with  $\text{dom } g = \bar{\mathbb{R}}_+^d$  such that*

$$\int_{\bar{\mathbb{R}}_+^d} h \circ g dx = C < \infty.$$

*Then the following are true:*

1. *For a sublevel set  $\text{lev}_y g$  with  $y > y_0$  we have:*

$$\mu[(\text{lev}_y g)^c] \leq C/h(y).$$

2. *For any point  $x_0 \in \mathbb{R}_+^d$  and any subgradient  $a \in \partial g(x_0)$  all coordinates of  $a$  are nonpositive. If in addition  $g(x_0) > y_0$  then all coordinates of  $a$  are negative.*
3. *For any point  $x_0 \in \bar{\mathbb{R}}_+^d$  such that  $g(x_0) > y_0$  we have:*

$$h \circ g(x_0) \leq \frac{Cd!}{d^d |x_0|},$$

*where  $|x| \equiv \prod_{k=1}^d x_k$  for  $x \in \mathbb{R}_+^d$ .*

4. *The function  $h$  reverses partial order on  $\bar{\mathbb{R}}_+^d$ : if  $x_1 < x_2$  then  $g(x_1) \geq g(x_2)$  and the last inequality is strict if  $g(x_1) > y_0$ .*
5. *The supremum of  $g$  on  $\bar{\mathbb{R}}_+^d$  is attained at 0.*

PROOF. 1. Since  $h$  is nondecreasing we have  $h(y) > 0$  and:

$$C = \int_{\bar{\mathbb{R}}_+^d} h \circ g dx \geq \int_{(\text{lev}_y g)^c} h \circ g dx \geq h(y) \mu[(\text{lev}_y g)^c].$$



2. Consider the linear function  $l(x) = a^T(x - x_0) + g(x_0)$ . We have  $g \geq l$ . If the vector  $a$  has a nonnegative coordinate  $a_i$  then consider a closed ball  $B = \bar{B}(x_0) \subset \mathbb{R}_+^d$ . If  $m$  is a minimum of the function  $l$  on  $B$  then the minimum of the function  $h \circ l$  on  $B + \lambda e_i$  is equal to  $h(m + \lambda a_i)$ , where  $e_i$  is the element of the basis which corresponds to the  $i$ th coordinate. For  $\lambda > 0$  we have  $B + \lambda e_i \subset \mathbb{R}_+^d$ .  
If  $a_i > 0$  then:

$$\int_{\mathbb{R}_+^p} h \circ g dx \geq \int_{\mathbb{R}_+^p} h \circ l dx \geq \int_B h \circ l dx \geq \mu[B] h(m + \lambda a_i) \rightarrow +\infty$$

as  $\lambda \rightarrow \infty$ , which contradicts the assumption.

If  $a_i = 0$  and  $g(x_0) = l(x_0) > y_0$ , then we can choose the radius of the ball small enough so that  $m > y_0$ . Then:

$$\int_{\mathbb{R}_+^p} h \circ g dx \geq \int_{\mathbb{R}_+^p} h \circ l dx \geq \int_K h \circ l dx \geq \mu[K] h(m) = +\infty$$

where  $K \equiv \cup_{\lambda > 0} (B + \lambda e_i)$ , and this again contradicts the assumption.

3. Consider the subgradient  $a \in \partial g(x_0)$ . For the linear function  $l(x) = a^T(x - x_0) + g(x_0)$  we have  $g \geq l$  and  $l(x_0) = g(x_0)$  therefore  $(\text{lev}_{g(x_0)} l)^c \subseteq (\text{lev}_{g(x_0)} g)^c$ . From the previous statement we have that  $\text{lev}_{g(x_0)} l$  is a simplex and using the Cauchy-Schwartz inequality we have:

$$\mu[(\text{lev}_{g(x_0)} l)^c] = \frac{(a^T x_0)^d}{d! |a|} \geq \frac{d^d |x_0|}{d!}$$

which together with 1. proves the statement.

4. Since  $x_1 \in \overline{\mathbb{R}_+^d}$  and  $x_1 < x_2$  we have  $x_2 \in \mathbb{R}_+^d = \text{ri}(\text{dom } g)$ . For any subgradient  $a \in \partial g(x_2)$  we have

$$g(x_1) - g(x_2) \geq a^T(x_1 - x_2) \geq 0$$

from the previous statement. Now, if  $g(x_1) > y_0$  then we can assume that  $g(x_2) > y_0$  since otherwise the statement is trivial. In this case all coordinates of  $a$  are negative and:

$$g(x_1) - g(x_2) \geq a^T(x_1 - x_2) > 0.$$

5. From the previous statement we have that  $h \circ g \leq h \circ g(0)$  on  $\mathbb{R}_+^d$  which together with continuity of  $h \circ g$  implies the statement.  $\square$

LEMMA 3.2. *Let  $h$  be an increasing transformation,  $g$  be a closed proper convex function on  $\overline{\mathbb{R}}_+^d$  and  $Q$  be a  $\sigma$ -finite Borel measure on  $\overline{\mathbb{R}}_+^d$ . Then:*

$$\int_{\text{lev}_a g} h \circ g dQ = \int_{-\infty}^a h'(y) Q[(\text{lev}_y g)^c \cap \text{lev}_a g] dy.$$

PROOF. Using the Fubini-Tonelli theorem we have:

$$\begin{aligned} \int_{\text{lev}_a g} h \circ g dQ &= \int_{\text{lev}_a g} \int_0^{h(a)} 1\{z \leq h \circ g(x)\} dz dQ(x) \\ &= \int_{\text{lev}_a g} \int_0^{h(a)} 1\{h^{-1}(z) \leq g(x)\} dz dQ(x) \\ &= \int_{\text{lev}_a g} \int_{-\infty}^a h'(y) 1\{y \leq g(x)\} dy dQ(x) \\ &= \int_{-\infty}^a h'(y) \int_{\text{lev}_a g} 1\{y \leq g(x)\} dQ(x) dy \\ &= \int_{-\infty}^a h'(y) Q[(\text{lev}_y g)^c \cap \text{lev}_a g] dy. \end{aligned}$$

□

LEMMA 3.3. *Let  $h$  be an increasing transformation and let  $g$  be a polyhedral convex function with  $\text{dom } g = \overline{\mathbb{R}}_+^d$  such that:*

$$\int_{\overline{\mathbb{R}}_+^d} h \circ g dx < \infty.$$

*Then  $g(0) < y_\infty$ .*

PROOF. For  $y_\infty = +\infty$  the statement is trivial so we assume that  $y_\infty$  is finite. If  $g(0) > y_\infty$  then since  $g$  is continuous there exists a ball  $B \subset \overline{\mathbb{R}}_+^d$  small enough such that  $g > y_\infty$  on  $B$  and therefore

$$\int_{\overline{\mathbb{R}}_+^d} h \circ g dx = \infty.$$

Let us assume that  $g(0) = y_\infty$ . By Lemma 4.13 there exists  $a \in \partial g(0)$  and therefore  $g(x) \geq l(x) \equiv a^T x + y_\infty$ . Let  $a_m$  be the minimum among the coordinates of the vector  $a$  and  $-1$ . Then on  $\overline{\mathbb{R}}_+^d$  we have  $l(x) \geq l_1(x) \equiv a_m \mathbf{1}^T x + y_\infty$  where  $a_m < 0$  and thus  $l_1(x) \leq y_\infty$ . By Lemma 3.2 we have:

$$\int_{\overline{\mathbb{R}}_+^d} h \circ g dx \geq \int_{\overline{\mathbb{R}}_+^d} h \circ l_1 dx = \int_{-\infty}^{y_\infty} h'(y) \mu[(\text{lev}_y l_1)^c \cap \overline{\mathbb{R}}_+^d] dy.$$

The set  $A_y = (\text{lev}_y g)^c \cap \overline{\mathbb{R}}_+^d$  is a simplex and:

$$\mu[A_y] = \frac{(y_\infty - y)^d}{d!(-a_m)^d}$$

for  $y \leq y_\infty$ . By assumption I.2 we have  $h'(y) \asymp (y_\infty - y)^{-\beta-1}$  as  $y \uparrow y_\infty$  where  $\beta > d$  and therefore:

$$\int_{\overline{\mathbb{R}}_+^d} h \circ g_1 dx = \int_{\overline{\mathbb{R}}_+^d} h \circ g dx = +\infty.$$

This contradiction proves that  $g(0) < y_\infty$ .  $\square$

LEMMA 3.4. *Let  $h$  be an increasing transformation and let  $l(x) = a^T x + b$  be a linear function such that all coordinates of  $a$  are negative and  $b < y_\infty$ . Then:*

$$\int_{\overline{\mathbb{R}}_+^d} h \circ l dx < \infty.$$

PROOF. We have  $l \leq b$  on  $\overline{\mathbb{R}}_+^d$  and by Lemma 3.2:

$$\int_{\overline{\mathbb{R}}_+^d} h \circ l dx = \int_{-\infty}^b h'(y) \mu[(\text{lev}_y l)^c \cap \overline{\mathbb{R}}_+^d] dy.$$

The set  $A_y = (\text{lev}_y l)^c \cap \overline{\mathbb{R}}_+^d$  is a simplex and:

$$\mu[A_y] = \frac{(b - y)^d}{d!|-a|}$$

for  $y \leq b$ . By assumption I.1 we have  $h'(y) = o(y^{-\alpha-1})$  as  $y \rightarrow -\infty$  for  $\alpha > d$  and therefore the integral is finite.  $\square$

LEMMA 3.5. *Let  $h$  be an increasing transformation and suppose that  $K \subset \overline{\mathbb{R}}_+^d$  is a compact set. Then there exists a closed proper convex function  $g \in \mathcal{G}(h)$  such that  $g > y_0$  on  $K$ .*

PROOF. If  $y_0 = -\infty$  then consider the function  $T(c)$  defined as:

$$T(c) = \int_{\overline{\mathbb{R}}_+^d} h \circ (-\mathbf{1}^T x + c) dx.$$

By Lemma 3.4,  $T(c)$  is finite for  $c < y_\infty$ , and by Lemma 3.3, we conclude that  $T(y_\infty) = +\infty$ . By monotone convergence  $T$  is left-continuous

for  $c \in (-\infty, y_\infty]$  and by dominated convergence is right-continuous for  $c \in (-\infty, y_\infty)$ . Since  $T(-\infty) = 0$  there exists  $c_1 < y_\infty$  such that  $T(c_1) = 1$  and thus the linear function  $l(x) = -\mathbf{1}^T x + c_1$  belongs to  $\mathcal{G}(h)$ .

If  $y_0 < -\infty$  then choose  $M$  such that  $\mathbf{1}^T x < M$  on  $K$ . Consider the function  $T(c)$  defined as:

$$T(c) = \int_{\mathbb{R}_+^d} h \circ (c(-\mathbf{1}^T x + M) + y_0) dx.$$

By Lemma 3.4,  $T(c)$  is finite for  $c < (y_\infty - y_0)/M$  and by Lemma 3.3,  $T((y_\infty - y_0)/M) = +\infty$ . By monotone and dominated convergence  $T$  is continuous for  $c \in [0, (y_\infty - y_0)/M]$ . Since  $T(0) = 0$  there exists  $c_1 \in (0, (y_\infty - y_0)/M)$  such that linear function  $l(x) = c_1(-\mathbf{1}^T x + M) + y_0$  belongs to  $\mathcal{G}(h)$ . By construction  $l > y_0$  on  $K$ .  $\square$

### 3.2. Preliminaries: properties of decreasing transformations.

LEMMA 3.6. *Let  $h$  be a decreasing transformation and  $g$  be a closed proper convex function such that*

$$\int_{\mathbb{R}^d} h \circ g dx = C < \infty.$$

*Then the following are true:*

1. *For  $y < +\infty$  the sublevel sets  $\text{lev}_y g$  are bounded and we have:*

$$\mu[\text{lev}_y g] \leq C/h(y);$$

2. *The infimum of  $g$  is attained at some point  $x \in \mathbb{R}^d$ .*

PROOF. 1. We have:

$$\begin{aligned} C &= \int_{\mathbb{R}_+^d} h \circ g dx \geq \int_{\text{lev}_y g} h \circ g dx \geq h(y) \mu[\text{lev}_y g] \\ \mu[\text{lev}_y g] &\leq C/h(y). \end{aligned}$$

The sublevel set  $\text{lev}_y g$  has the same dimension as  $\text{dom } g$  (Theorem 7.6 Rockafellar [1970]) which is  $d$ . By Lemma 4.1 this set is bounded when  $y < y_0$ . Therefore, it is enough to prove that  $\text{lev}_{y_0} g$  is bounded for  $y_0 < +\infty$ .

Since  $h \circ g$  is a density we have  $\inf g < y_0$ . If  $g$  is constant on  $\text{dom } g$ , then for all  $y \in [\inf g, +\infty)$  we have  $\text{lev}_y g = \text{lev}_{\inf g} h$  and is therefore bounded. Otherwise, we can choose  $\inf h \leq y_1 < y_2 < y_0$ . Then  $\mu[\text{lev}_{y_2} g] < \infty$ , and by Lemma 4.3 we have  $\mu[\text{lev}_{y_0} g] < \infty$ . The argument above shows that  $\text{lev}_{y_0} g$  is also bounded.

2. Follows from the fact that  $g$  is continuous and  $\text{lev}_y g$  is bounded and nonempty for  $y > \inf g$ .  $\square$

LEMMA 3.7. *Let  $h$  be a decreasing transformation, let  $g$  be a closed proper convex function on  $\mathbb{R}^d$ , and let  $Q$  be a  $\sigma$ -finite Borel measure on  $\mathbb{R}^d$ . Then:*

$$\int_{(\text{lev}_a g)^c} h \circ g dQ = - \int_a^{+\infty} h'(y) Q[\text{lev}_y g \cap (\text{lev}_a g)^c] dy.$$

PROOF. Using the Fubini-Tonelli theorem we have:

$$\begin{aligned} \int_{(\text{lev}_a g)^c} h \circ g dQ &= \int_{(\text{lev}_a g)^c} \int_0^{h(a)} 1\{z \leq h \circ g(x)\} dz dQ(x) \\ &= \int_{(\text{lev}_a g)^c} \int_0^{h(a)} 1\{h^{-1}(z) \geq g(x)\} dz dQ(x) \\ &= - \int_{(\text{lev}_a g)^c} \int_a^{+\infty} h'(y) 1\{y \geq g(x)\} dy dQ(x) \\ &= - \int_a^{+\infty} h'(y) \int_{(\text{lev}_a g)^c} 1\{y \geq g(x)\} dQ(x) dy \\ &= - \int_a^{+\infty} h'(y) Q[\text{lev}_y g \cap (\text{lev}_a g)^c] dy. \end{aligned}$$

$\square$

LEMMA 3.8. *Let  $h$  be an decreasing transformation and let  $g$  be a closed proper convex function such that:*

$$\int_{\mathbb{R}^d} h \circ g dx < \infty.$$

*Then  $\inf g > y_\infty$ .*

PROOF. Since  $g$  is proper the statement is trivial for  $y_\infty = -\infty$ , so we assume that  $y_\infty > -\infty$ . If for  $x_0$  we have  $g(x_0) = y_\infty$ , then there exists a ball  $B \equiv B(x; r)$  such that  $g < y_\infty + \varepsilon$  on  $B$ . Consider the convex function  $f$  defined as:

$$f(x) = y_\infty + (\varepsilon/r) \|x - x_0\| + \delta(x \mid B).$$

Then by convexity  $f \geq g$  and:

$$\int_{\mathbb{R}^d} h \circ g dx \geq \int_{\mathbb{R}^d} h \circ f dx.$$

We have  $\mu[\text{lev}_y f] = S(y - y_\infty)^d$  for  $y \in [y_\infty, y_\infty + \varepsilon]$  where  $S$  is the Lebesgue measure of a unit ball  $B(0; 1)$  and by Lemma 3.7 we can compute:

$$\int_{\mathbb{R}^d} h \circ f dx = -S \int_{y_\infty}^{y_\infty + \varepsilon} h'(y)(y - y_\infty)^d dy.$$

The assumption D.2 implies:

$$\int_{\mathbb{R}^d} h \circ g dx \geq \int_{\mathbb{R}^d} h \circ f dx = \infty,$$

which proves the statement.  $\square$

LEMMA 3.9. *Let  $h$  be a decreasing transformation. Then for any convex function  $g$  such that  $h \circ g$  belongs to the decreasing model  $\mathcal{P}(h)$  we have:*

$$\int_{\mathbb{R}^d} [h \log h] \circ g dx < \infty.$$

PROOF. By assumption D.1 the function  $-[h \log h](y)$  is decreasing to zero as  $y \rightarrow +\infty$  and we have:

$$0 < -[h \log h](y) < Cy^{-d-\alpha'}$$

for  $C$  large enough and  $\alpha' \in (0, \alpha)$  as  $y \rightarrow +\infty$ .

By Lemma 3.6 the level sets  $\text{lev}_y g$  are bounded and since  $h \circ g \in \mathcal{P}(h)$  we have  $\inf g > y_\infty$ . Therefore, the integral is finite if and only if the integral:

$$\int_{(\text{lev}_a g)^c} [h \log h] \circ g dx$$

is finite for some  $a > y_\infty$ . Choosing  $a$  large enough and using Lemma 3.7 for the decreasing transformation  $h_1(y) = y^{-d-\alpha'}$  we obtain:

$$\begin{aligned} 0 &\geq \int_{(\text{lev}_a g)^c} [h \log h] \circ g dx \geq -C \int_{(\text{lev}_a g)^c} h_1 \circ g dx \geq C \int_a^{+\infty} h'_1(y) \mu[\text{lev}_y g] dy \\ &= -C(d + \alpha') \int_a^{+\infty} y^{-d-\alpha'-1} \mu[\text{lev}_y g] dy. \end{aligned}$$

By Lemma 4.3 we have  $\mu(\text{lev}_y g) = O(y^d)$  and therefore the last integral is finite.  $\square$

LEMMA 3.10. *Let  $h$  be a decreasing transformation and suppose that  $K \subset \mathbb{R}^d$  is a compact set. Then there exists a closed proper convex function  $g \in \mathcal{G}(h)$  such that  $g < y_0$  on  $K$ .*

PROOF. Let  $B$  be a ball such that  $K \subset B$ . Let  $c$  be such that  $h(c) = 1/\mu[B]$ . Then the function  $g \equiv c + \delta(\cdot | B)$  belongs to  $\mathcal{G}(h)$ .  $\square$

3.3. *Proofs for Existence Results.* Before giving proofs of Theorems 2.13 and 2.14, we establish two auxiliary lemmas.

A set of points  $x = \{x_i\}_{i=1}^n$  in  $\mathbb{R}^d$  is in *general position* if for any subset  $x' \subseteq x$  of the size  $d + 1$  the Lebesgue measure of  $\text{conv}(x')$  is not zero.

LEMMA 3.11. *If  $X_1, \dots, X_n$  are i.i.d.  $p_0 = h \circ g_0 \in \mathcal{P}(h)$  for a monotone transformation  $h$ , then the observations  $X$  are in general position with probability 1.*

PROOF. Points are not in general position if at least one subset  $Y$  of  $X$  of size  $d + 1$  belongs to a proper linear subspace of  $\mathbb{R}^d$ . This is true if and only if  $X$  as a vector in  $\mathbb{R}^{nd}$  belongs to a certain non-degenerate algebraic variety. Since with probability 1 we have  $X \subset \text{dom } g_0$  and by definition  $\dim(\text{dom } g_0) = d$ , the statement follows from Okamoto [1973].  $\square$

Below we assume that our observations are in general position for any  $n$ . For an increasing model we also assume that all  $X_i$  belong to  $\mathbb{R}_+^d$ . This assumption holds with probability 1 since  $\mu[\mathbb{R}_+^d \setminus \mathbb{R}_+^d] = 0$ .

If a MLE for the model  $\mathcal{P}(h)$  exists, then it maximizes the functional:

$$\mathbb{L}_n g \equiv \int (\log h) \circ g d\mathbb{P}_n$$

over  $g \in \mathcal{G}(h)$ , where the last integral is over  $\mathbb{R}_+^d$  for increasing  $h$  and over  $\mathbb{R}^d$  for decreasing models. The theorem below determines the form of the MLE for an increasing model. We write  $\text{ev}_x f = (f(x_1), \dots, f(x_n))$ ,  $x = (x_1, \dots, x_n)$  with  $x_i \in \mathbb{R}^d$ .

LEMMA 3.12. *Consider an increasing transformation  $h$ . For any convex function  $g$  with  $\text{dom } g = \mathbb{R}_+^d$  such that:  $\int_{\mathbb{R}_+^d} h \circ g dx \leq 1$  and  $\mathbb{L}_n g > -\infty$ , there exists  $\tilde{g} \in \mathcal{G}(h)$  such that  $\tilde{g} \geq g$  and  $\mathbb{L}_n \tilde{g} \geq \mathbb{L}_n g$ . The function  $\tilde{g}$  can be chosen as a minimal element in  $\text{ev}_X^{-1} \tilde{p}$  where  $\tilde{p} = \text{ev}_X g$ .*

PROOF. Let  $p = \text{ev}_X g$ . Since  $\mathbb{L}_n g > -\infty$  we have  $g(X_i) > y_0$  for all  $1 \leq i \leq n$  and therefore  $g > y_0$  for  $x \in \text{conv}(X)$ . Consider any minimal element  $g_1$  among convex functions in  $\text{ev}_X^{-1} p$  (which exists by Lemma 4.15). Then:

$$\int_{\mathbb{R}_+^d} h \circ g_1 dx \leq \int_{\mathbb{R}_+^d} h \circ g dx \leq 1.$$

Since  $g_1$  is polyhedral we have  $g_1 = \max l_i$  for some linear functions  $l_i(x) = a_i^T x + b$  and for each function  $l_i$  there exists some facet of  $g_1$  such that  $g_1 = l_i$  on it.

By Lemma 4.15 the interior of the facet of  $g_1$  which corresponds to  $l_i$  contains some  $X_{j_i} \in X$ . We have  $\partial g_1(X_{j_i}) = \{a_i\}$  and  $g_1(X_{j_i}) = g(X_{j_i}) > y_0$ . Thus by Lemma 3.1, all coordinates of  $a_i$  are negative and the supremum  $M$  of  $g_1$  is attained at 0. Therefore  $b_i = l_i(0) \leq M$ . By Lemma 3.3 we have  $M < y_\infty$ . Thus by Lemma 3.4 the functions  $h \circ (l_i + c)$  are integrable for all  $c < y_\infty - M$ . Since  $g_1$  has only finite number of facets we have that  $h \circ (g_1 + c)$  is also integrable for all  $c < y_\infty - M$ . Finally, for  $c = y_\infty - M$  function  $h \circ (g_1 + c)$  is not integrable by Lemma 3.3.

The function  $T(c)$  defined as:

$$T(c) \equiv \int_{\mathbb{R}_+^d} h \circ (g_1 + c) dx$$

is increasing, finite for  $c \in [0, y_\infty - M)$  and continuous for  $c \in [0, y_\infty - M]$  by monotone and dominated convergence. Since  $T(0) \leq 1$  and  $T(y_\infty - M) = +\infty$ , there exists  $c_1 \in (0, y_\infty - M)$  such that  $T(c_1) = 1$ . Then the function  $\tilde{g} \equiv g_1 + c_1$  satisfies the conditions of our lemma.  $\square$

**THEOREM 3.13.** *If an MLE  $\hat{g}_0$  exists for the increasing model  $\mathcal{P}(h)$ , then there exists an MLE  $\hat{g}_1$  which is a minimal element in  $\text{ev}_X^{-1} p$  where  $q = \text{ev}_X \hat{g}_0$ . In other words  $\hat{g}_1$  is a polyhedral convex function such that  $\text{dom } g_1 = \mathbb{R}_+^d$ , and the interior of each facet contains at least one element of  $X$ . If  $h$  is strictly increasing on  $[y_0, y_\infty]$ , then  $\hat{g}_0(x) = \hat{g}_1(x)$  for all  $x$  such that  $\hat{g}_0(x) > y_0$  and thus defines the same density from  $\mathcal{P}(h)$ .*

**PROOF.** Let  $\hat{g}_0$  be any MLE. Then by Lemma 3.5 applied to  $K = \text{conv}(X)$  it follows that  $\mathbb{L}_n \hat{g}_0 > -\infty$ . By Lemma 3.12 there exists a function  $\hat{g}_1 \in \mathcal{P}(h)$  such that  $\hat{g}_1$  is a minimal element in  $\text{ev}_X^{-1} q_1$  where  $q_1 = \text{ev}_X \hat{g}_1$  and  $\hat{g}_1 \geq \hat{g}_0$ . Since  $\hat{g}_0$  is a MLE we have  $\text{ev}_X \hat{g}_0 = \text{ev}_X \hat{g}_1$  which together with Lemma 4.15 proves the first part of the statement.

By Lemma 3.3 we have  $\hat{g}_0 < y_\infty$  and  $\hat{g}_1 < y_\infty$ . Since  $h \circ \hat{g}_0$  and  $h \circ \hat{g}_1$  are continuous functions, for the strictly increasing  $h$  the equality:

$$\int_{\mathbb{R}_+^d} (h \circ \hat{g}_1 - h \circ \hat{g}_0) dx = 0$$

implies that  $\hat{g}_1(x) = \hat{g}_0(x)$  for  $x$  such that  $\hat{g}_0(x) > y_0$ .  $\square$

Here are the corresponding results for decreasing transformations  $h$ .

**LEMMA 3.14.** *Consider a decreasing transformation  $h$ . For any convex function  $g$  such that:*

$$\int_{\mathbb{R}^d} h \circ g dx \leq 1$$



and  $\mathbb{L}_n g > -\infty$  there exists  $\tilde{g} \in \mathcal{G}(h)$  such that  $\tilde{g} \leq g$  and  $\mathbb{L}_n \tilde{g} \geq \mathbb{L}_n g$ . The function  $\tilde{g}$  can be chosen as the maximal element in  $\text{ev}_X^{-1} \tilde{q}$  where  $\tilde{q} = \text{ev}_X \tilde{g}$ .

PROOF. Let  $q = \text{ev}_X g$ . Since  $\mathbb{L}_n g > 0$  we have  $g(X_i) < y_0$  for all  $1 \leq i \leq n$  and therefore  $g < y_0$  for  $x \in \text{conv}(X)$ . Consider the maximal element  $g_1$  among convex functions in  $\text{ev}_X^{-1} q$  (which exists and is unique by Lemma 4.14). Then:

$$\int_{\mathbb{R}^d} h \circ g_1 dx \leq \int_{\mathbb{R}_+^d} h \circ g dx = 1.$$

By Lemma 3.6 there exists  $x_0$  and  $m > -\infty$  such that  $g \geq g(x_0) = m$ . By Lemma 3.8 we have  $m > y_\infty$ . By Lemma 4.14 we have  $\text{dom } g_1 = \text{conv}(X)$  and therefore:

$$\int_{\mathbb{R}^d} h \circ (g_1 + c) dx \leq h(m) \mu[\text{conv}(X)] < \infty.$$

The function  $T(c)$  defined as:

$$T(c) \equiv \int_{\mathbb{R}_+^d} h \circ (g_1 + c) dx$$

is increasing, finite for  $c \in (y_\infty - m, 0]$  and continuous for  $c \in [y_\infty - m, 0]$  by monotone and dominated convergence. Since  $T(0) \leq 1$  and  $T(y_\infty - m) = +\infty$ , there exists  $c_1 \in (y_\infty - m, 0)$  such that  $T(c_1) = 1$ . Then the function  $\tilde{g} \equiv g_1 + c_1$  satisfies the conditions of our lemma.  $\square$

**THEOREM 3.15.** *If the MLE  $\hat{g}_0$  exists for the decreasing model  $\mathcal{P}(h)$ , then there exists another MLE  $\hat{g}_1$  which is the maximal element in  $\text{ev}_X^{-1} q$  where  $q = \text{ev}_X \tilde{g}_0$ . In other words  $\hat{g}_1$  is a polyhedral convex function with the set of knots  $K_n \subseteq X$  and domain  $\text{dom } \hat{g}_1 = \text{conv}(X)$ . If  $h$  is strictly decreasing on  $[y_\infty, y_0]$ , then  $\hat{g}_0(x) = \hat{g}_1(x)$ .*

PROOF. Let  $\hat{g}_0$  be any MLE. Then by Lemma 3.10 applied to  $K = \text{conv}(X)$  we have that  $\mathbb{L}_n \hat{g}_0 > 0$ . By Lemma 3.14 there exists a function  $\hat{g}_1 \in \mathcal{G}_h$  such that  $\hat{g}_1$  is the maximal element in  $\text{ev}_X^{-1} q_1$  where  $q_1 = \text{ev}_X \hat{g}_1$  and  $\hat{g}_1 \leq \hat{g}_0$ . Since  $\hat{g}_0$  is a MLE we have  $\text{ev}_X \hat{g}_0 = \text{ev}_X \hat{g}_1$ , which together with Lemma 4.14 proves the first part of the statement.

By Lemma 3.8 we have  $\hat{g}_0 \geq \hat{g}_1 > y_\infty$ . Since  $h \circ \hat{g}_0$  and  $h \circ \hat{g}_1$  are continuous functions, for the strictly increasing  $h$ , the equality:

$$\int_{\mathbb{R}_+^d} (h \circ \hat{g}_1 - h \circ \hat{g}_0) dx = 0$$

implies that  $\widehat{g}_1(x) = \widehat{g}_0(x)$  for  $x \in \text{conv}(X)$ . Therefore  $\widehat{g}_0(x) \geq y_\infty$  for  $x \notin \text{conv}(X)$ . Since  $\widehat{g}_0$  is convex we have  $\widehat{g}_0 = \widehat{g}_1$ .  $\square$

The bounds provided by the following key lemma are the remaining preparatory work for proving existence of the MLE in the case of increasing transformations.

For an increasing model  $\mathcal{P}(h)$  let us denote by  $\mathcal{N}(h, X, \varepsilon)$  for  $\varepsilon > -\infty$  the family of all convex functions  $g \in \mathcal{G}(h)$  such that  $g$  is a minimal element in  $\text{ev}_X^{-1} q$  where  $q = \text{ev}_X g$  and  $\mathbb{L}_n g \geq \varepsilon$ . By Lemma 3.5, the family  $\mathcal{N}(h, X, \varepsilon)$  is not empty for  $\varepsilon > -\infty$  small enough. By construction for  $g \in \mathcal{N}(h, X, \varepsilon)$  we have  $g(X_i) > y_0$  for  $X_i \in X$ .

LEMMA 3.16. *There exist constants  $c(x, X, \varepsilon)$  and  $C(x, X, \varepsilon) < y_\infty$  which depend only on  $x \in \mathbb{R}_+^d$ , the observations  $X$ , and  $\varepsilon$ , such that for any  $g \in \mathcal{N}(h, X, \varepsilon)$  we have:*

$$c(x, X, \varepsilon) \leq g(x) \leq C(x, X, \varepsilon).$$

PROOF. By Lemma 3.1 we have:

$$h \circ g(X_i) \leq \frac{d!}{d^d |X_i|},$$

which gives the upper bounds  $C(X_i, X, \varepsilon)$ . By assumption we have:

$$(\max h \circ g(X_i))^{n-1} \min h \circ g(X_i) \geq \prod h \circ g(X_i) \geq e^{n\varepsilon},$$

and therefore:

$$\min h \circ g(X_i) \geq \frac{e^{n\varepsilon}}{h(\max C(X_i, X, \varepsilon))^{n-1}},$$

which gives the uniform lower bound  $c(X_i, X, \varepsilon)$  for all  $X_i \in X$ . Since by Lemma 3.1  $g(0) \geq g(X_i)$  we also obtain  $c(0, X, \varepsilon)$ .

Now, we prove that there exist  $C(0, X, \varepsilon)$ . Let  $l$  be a linear function which defines any facet of  $g$  for which 0 is an element. By Lemma 4.15, there exists  $X_a \in X$  which belongs to this facet. Then  $g(0) = l(0)$  and  $g(X_a) = l(X_a)$ . Let us denote by  $S$  the simplex  $\{l = l(X_a)\} \cap \mathbb{R}_+^d$ , by  $S^*$  the simplex  $\{l \geq l(X_a)\} \cap \mathbb{R}_+^d$  and by  $l'$  the linear function which is equal to  $c \equiv \min c(X_i, X, \varepsilon)$  on  $S$  and to  $g(0)$  at 0. By the Cauchy-Schwartz inequality (as in the proof of Lemma 3.1) we have:

$$\mu[S^*] \geq \frac{d^d |X_a|}{d!}$$

We have  $l \geq l'$ :

$$1 = \int_{\overline{\mathbb{R}}_+^d} h \circ g dx \geq \int_{S^*} h \circ l' dx.$$

By Lemma 3.2

$$\begin{aligned} \int_{S^*} h \circ l' dx &= \mu[S^*] \int_c^{g(0)} h'(y) \left( \frac{g(0) - y}{g(0) - c} \right)^d dy \\ &\geq \frac{d^d |X_a|}{d!} \int_c^{y_\infty} h'(y) 1\{y \leq g(0)\} \left( \frac{g(0) - y}{g(0) - c} \right)^d dy. \end{aligned}$$

Consider the function  $T(s)$  defined as:

$$T(s) = \frac{d^d |X_a|}{d!} \int_c^{y_\infty} h'(y) 1\{y \leq s\} \left( \frac{s - y}{s - c} \right)^d dy.$$

If  $y_\infty = +\infty$ , then for a fixed  $y \in (c, +\infty)$  we have:

$$h'(y) 1\{y \leq s\} \left( \frac{s - y}{s - c} \right)^d \uparrow h'(y) \text{ as } s \rightarrow y_\infty$$

and by monotone convergence we have:

$$T(s) \uparrow \int_c^{y_\infty} h'(y) dy = +\infty \text{ as } s \rightarrow y_\infty.$$

If  $y_\infty < +\infty$  then for a fixed  $y \in (c, y_\infty]$  we have:

$$h'(y) 1\{y \leq s\} \left( \frac{s - y}{s - c} \right)^d \uparrow h'(y) \left( \frac{y_\infty - y}{y_\infty - c} \right)^d \text{ as } s \rightarrow y_\infty$$

and by monotone convergence we have:

$$T(s) \uparrow \int_c^{y_\infty} h'(y) \left( \frac{y_\infty - y}{y_\infty - c} \right)^d dy = +\infty \text{ as } s \rightarrow y_\infty,$$

by assumption I.2.

Thus there exists  $s_0 \in (c, y_\infty)$  such that  $T(s_0) > 1$ . This implies  $g(0) < s_0$ . Since  $s_0$  depends only on  $X_a$  and  $\min c(X_i, X, \varepsilon)$  this gives an upper bound  $C(0, X, \varepsilon)$ .

By Lemma 3.1 for any  $x_0 \in \overline{\mathbb{R}}_+^d$  we can set  $C(x_0, X, \varepsilon) = C(0, X, \varepsilon)$ . Let  $l(x) = a^T x + l(0)$  be a linear function which defines the facet of  $g$  to which  $x$  belongs. By Lemma 4.15 there exists  $X_a \in X$  which belongs to this facet,

and thus  $l(X_a) = g(X_a)$ . By Lemma 3.1 we have  $a_k < 0$  for all  $k$ , and by definition  $l(0) \leq g(0)$ . We have

$$c(X_a, X, \varepsilon) \leq g(X_a) = l(X_a) = a^T X_a + l(0) \leq a^T X_a + g(0),$$

and therefore

$$\begin{aligned} a_k &\geq \frac{c(X_a, X, \varepsilon) - C(0, X, \varepsilon)}{(X_a)_k} \\ l(0) &\geq c(X_a, X, \varepsilon). \end{aligned}$$

Now,

$$g(x_0) = l(x_0) \geq \frac{c(X_a, X, \varepsilon) - C(0, X, \varepsilon)}{(X_a)_k} (x_0)_k + c(X_a, X, \varepsilon).$$

Since we have only finite number of possible choices for  $X_a$ , we obtained  $c(x_0, X, \varepsilon)$  which concludes the proof.  $\square$

Now we are ready for the proof of Theorem 2.13.

PROOF. (Theorem 2.13) By Lemma 3.5 there exists  $\varepsilon$  small enough such that the family  $\mathcal{N}(h, X, \varepsilon)$  is not empty. Clearly, we can restrict MLE candidates  $\hat{g}$  to functions in the family  $\mathcal{N}(h, X, \varepsilon)$ . The set  $N = \text{ev}_X \mathcal{N}(h, X, \varepsilon)$  is bounded by Lemma 3.16. Let us denote by  $q^*$  a point in the closure  $\bar{N}$  of  $N$  which maximizes the continuous function:

$$L_n(p) = \frac{1}{n} \sum_{i=1}^n \log h(q_i).$$

Since  $q \in \bar{N}$ , there exists a sequence of functions  $g_k \in \mathcal{N}(h, X, \varepsilon)$  such that  $\text{ev}_X g_k$  converges to  $q^*$ . By Theorem 10.9 Rockafellar [1970] and Lemma 3.16 there exists a finite convex function  $g^*$  on  $\bar{\mathbb{R}}_+^d$  such that some subsequence  $g_l$  pointwise converges to  $g^*$ . Therefore we have  $\text{ev}_X g^* = q^*$ . Since  $X \subset \mathbb{R}_+^d$  we can assume that  $g^*$  is closed. By Fatou's lemma we have:

$$\int_{\mathbb{R}_+^d} h \circ g^* dx \leq 1.$$

By Lemma 3.12 there exists  $g \in \mathcal{G}(h)$  such that  $g \geq g^*$  and  $\mathbb{L}_n g \geq \mathbb{L}_n g^* = L_n(q^*)$ . By assumption this implies  $\mathbb{L}_n g = \mathbb{L}_n g^*$ . Hence  $g$  is the MLE. Finally, we have to add the almost surely clause since we assumed that the points  $X_i$  belong to  $\mathbb{R}_+^d$ .  $\square$

Before proving existence of the MLE for a decreasing transformation family, we need two lemmas.

LEMMA 3.17. *Consider a decreasing model  $\mathcal{P}(h)$ . Let  $\{g_k\}$  be a sequence of convex functions from  $\mathcal{G}(h)$ , and let  $\{n_k\}$  be a nondecreasing sequence of positive integers  $n_k \geq n_d$  such that for some  $\varepsilon > -\infty$  and  $\rho > 0$  the following is true:*

1.  $\mathbb{L}_{n_k} g_k \geq \varepsilon$ ;
2. if  $\mu[\text{lev}_{a_k} g_k] = \rho$  for some  $a_k$ , then  $\mathbb{P}_{n_k}[\text{lev}_{a_k} g_k] < d/n_d$ .

*Then there exists  $m > y_\infty$  such that  $g_k \geq m$  for all  $k$ .*

PROOF. Suppose, on the contrary, that  $m_k \rightarrow y_\infty$  where  $m_k = \min g_k$ . The first condition implies that  $X_d \equiv \{X_1, \dots, X_{n_d}\} \in \text{dom } h_k$ , and therefore by Corollary 4.4 the function  $\mu[\text{lev}_y g_k]$  as a function of  $y$  admits all values in the interval  $[\mu[\text{lev}_{m_k} g_k], \mu[\text{conv}(X_d)]]$ . If the second condition is true for some  $\rho$  then it is also true for all  $\rho' \in (0, \rho)$ , and therefore we can assume that  $\rho < \mu[\text{conv}(X_d)]$ .

By Lemma 3.6 we have  $\mu[\text{lev}_{m_k} g_k] \rightarrow 0$ , and thus there exists such  $a_k$  that  $\mu[\text{lev}_{a_k} g_k] = \rho$  for all  $k$  large enough. We define  $A_k = \text{lev}_{a_k} g_k$ . By Lemma 3.6 we have:  $h(a_k) \leq 1/\rho$  and therefore the sequence  $\{a_k\}$  is bounded below by some  $a > y_\infty$ .

Consider  $t_k > m_k$  such that  $t_k \rightarrow y_\infty$ . We will specify the exact form of  $t_k$  later in the proof. Since  $a_k$  are bounded away from  $y_\infty$ , it follows that for  $k$  large enough we will have  $t_k < a_k$ . Using Lemma 4.3 we obtain:

$$\rho = \mu[A_k] \leq \mu[\text{lev}_{t_k} g_k] \left[ \frac{a_k - m_k}{t_k - m_k} \right]^d \leq \frac{1}{h(t_k)} \left[ \frac{a_k - m_k}{t_k - m_k} \right]^d$$

which implies:

$$a_k \geq m_k + (t_k - m_k)[\rho h(t_k)]^{1/d}.$$

We have:

$$g_k \geq m_k 1\{A_k\} + a_k(1 - 1\{A_k\}),$$

and hence:

$$\begin{aligned} \mathbb{L}_{n_k} g_k &\leq \mathbb{P}_{n_k}(A_k) \log h(m_k) + (1 - \mathbb{P}_{n_k}(A_k)) \log h(a_k) \\ &\leq \mathbb{P}_{n_k}(A_k) \log h(m_k) + (1 - \mathbb{P}_{n_k}(A_k)) \log h(m_k + (t_k - m_k)[\rho h(t_k)]^{1/d}). \end{aligned}$$

*Case  $y_\infty = -\infty$ .* Choose  $t_k = (1 - \delta)m_k$  where  $\delta \in (0, 1)$ . Then starting from some  $k$  we have  $m_k < t_k$ ,  $h(m_k) > 1$ ,  $h(-Cm_k) < 1$  and  $\delta[\rho h(t_k)]^{1/d} > C + 1$ . This implies:

$$m_k + (t_k - m_k)[\rho h(t_k)]^{1/d} = m_k(1 - \delta[\rho h(t_k)]^{1/d}) \geq -Cm_k,$$

and hence:

$$\begin{aligned} \mathbb{L}_{n_k} g_k &\leq \mathbb{P}_{n_k}(A_k) \log h(m_k) + (1 - \mathbb{P}_{n_k}(A_k)) \log h(-Cm_k) \\ &\leq \frac{d}{n_d} \log h(m_k) + \frac{n_d - d}{n_d} \log h(-Cm_k) = \frac{d}{n_d} \log [h(m_k)h(-Cm_k)^\gamma] \rightarrow -\infty. \end{aligned}$$

*Case  $y_\infty > -\infty$ .* Without loss of generality we can assume that  $y_\infty = 0$ . Choose  $t_k = (1 + \delta)m_k$  where  $\delta > 0$ . Then:

$$m_k + (t_k - m_k)[\rho h(t_k)]^{1/d} \geq m_k \delta [\rho h((1 + \delta)m_k)]^{1/d} \asymp m_k^{-\frac{\beta-d}{d}} \rightarrow +\infty$$

which implies

$$h(m_k + (t_k - m_k)[\rho h(t_k)]^{1/d}) = o\left(m_k^{\frac{\alpha(\beta-d)}{d}}\right).$$

This in turn yields

$$\exp(\mathbb{L}_{n_k} g_k) = o\left(m_k^{-\frac{\beta d}{n_d} + \frac{\alpha(\beta-d)(n_d-d)}{d n_d}}\right) = o(1).$$

Therefore in both cases we obtained  $\mathbb{L}_{n_k} g_k \rightarrow -\infty$ . This contradiction concludes the proof.  $\square$

For a decreasing model  $\mathcal{P}(h)$ , let us denote by  $\mathcal{N}(h, X, \varepsilon)$  for  $\varepsilon > -\infty$  the family of all convex functions  $g \in \mathcal{G}(h)$  such that  $g$  is a maximal element in  $\text{ev}_X^{-1} q$  where  $q = \text{ev}_X g$ , and  $\mathbb{L}_n g \geq \varepsilon$ . By Lemma 3.10 family  $\mathcal{N}(h, X, \varepsilon)$  is not empty for  $\varepsilon > -\infty$  small enough. By construction for  $g \in \mathcal{N}(h, X, \varepsilon)$  we have  $g(X_i) < y_0$  for  $X_i \in X$ .

**LEMMA 3.18.** *For given observations  $X = (X_1, \dots, X_n)$  such that  $n \geq n_d$ , there exist constants  $m > y_\infty$  and  $M$  which depend only on observations  $X$  and  $\varepsilon$  such that for any  $g \in \mathcal{N}(h, X, \varepsilon)$  we have  $m \leq g(x) \leq M$  on  $\text{conv}(X)$ .*

**PROOF.** Since by assumption the points  $X$  are in general position, there exists  $\rho > 0$  such that for any  $d$ -dimensional simplex  $S$  with vertices from

$X$  we have  $\mu[S] \geq \rho$ . Then any convex set  $C \subseteq \text{conv}(X)$  such that  $\mu[C] = \rho$  cannot contain more than  $d$  points from  $X$ . Therefore, we have  $\mathbb{P}_n[C] \leq d/n \leq d/n_d$ .

An arbitrary sequence of functions  $\{g_k\}$  from  $\mathcal{N}(h, X, \varepsilon)$  satisfies the conditions of Lemma 3.17 with  $n_k \equiv n$ , the same  $\varepsilon$  and  $\rho$  constructed above. Therefore the sequence  $\{g_k\}$  is bounded below by some constant greater than  $y_\infty$ . Thus the family of functions  $\mathcal{N}(h, X, \varepsilon)$  is uniformly bounded below by some  $m > y_\infty$ .

Consider any  $g \in \mathcal{N}(h, X, \varepsilon)$ . Let  $M_g$  be the supremum of  $g$  on  $\text{dom } h$ . By Theorem 32.2 Rockafellar [1970], the supremum is obtained at some  $X_M \in X$  and therefore  $M_g < y_0$ . Let  $m_g$  be the minimum of  $g$  on  $X$ . We have:

$$\begin{aligned} h(m_g)^{n-1} h(M_g) &\geq e^{n\varepsilon} \\ h(M_g) &\geq \frac{e^{n\varepsilon}}{h(m_g)^{n-1}} \geq \frac{e^{n\varepsilon}}{h(m)^{n-1}}. \end{aligned}$$

Thus we obtained an upper bound  $M$  which depends only on  $m$  and  $X$ .  $\square$

Now we are ready for the proof of Theorem 2.14.

PROOF. (Theorem 2.14) By Lemma 3.10 there exists  $\varepsilon$  small enough such that the family  $\mathcal{N}(h, X, \varepsilon)$  is not empty. Clearly, we can restrict MLE candidates to the functions in the family  $\mathcal{N}(h, X, \varepsilon)$ . The set  $N = \text{ev}_X \mathcal{N}(h, X, \varepsilon)$  is bounded by Lemma 3.18. Let us denote by  $q^*$  the point in the closure  $\bar{N}$  of  $N$  which maximizes the continuous function:

$$L_n(q) = \frac{1}{n} \sum_{i=1}^n \log h(q_i).$$

Since  $q \in \bar{N}$ , there exists a sequence of functions  $g_k \in \mathcal{N}(h, X, \varepsilon)$  such that  $\text{ev}_X g_k$  converges to  $q^*$ . By Lemma 3.18 the functions  $f_k = \sup_{l \geq k} g_l$  are finite convex functions on  $\text{conv}(X)$ , the sequence  $\{f_k(x)\}$  is monotone decreasing for each  $x \in \text{conv}(X)$  and bounded below. Therefore  $f_k \downarrow g^*$  for some convex function  $g^*$ , and by construction  $\text{ev}_X g^* = q^*$ . We have:

$$\int_{\mathbb{R}_+^d} h \circ f_k dx \leq \int_{\mathbb{R}_+^d} h \circ g_k dx = 1$$

and thus by Fatou's lemma:

$$\int_{\mathbb{R}_+^d} h \circ g^* dx \leq 1.$$

By Lemma 3.14 there exists  $g \in \mathcal{G}(h)$  such that  $g \leq g^*$  and  $\mathbb{L}_n g \geq \mathbb{L}_n g^* = L_n(q^*)$ . By assumption this implies  $\mathbb{L}_n g = \mathbb{L}_n g^*$ . Thus the function  $g$  is the MLE.

Finally, we have to add the almost surely clause since we assumed that the points  $X_i$  are in general position.  $\square$

**3.4. Proofs for Consistency Results.** We begin with proofs for some technical results which we will use in the consistency arguments for both increasing and decreasing models. The main argument for proving Hellinger consistency proceeds along the lines of the proof given in the case of  $d = 1$  by Pal, Woodroffe and Meyer [2007].

**LEMMA 3.19.** *Consider a monotone model  $\mathcal{P}(h)$ . Suppose the true density  $h \circ g_0$  and the sequence of MLEs  $\{\hat{g}_n\}$  have the following properties:*

$$\int (h \log h) \circ g_0(x) dx < \infty,$$

and

$$\int \log[\varepsilon + h \circ \hat{g}_n(x)] d(\mathbb{P}_n(x) - P_0(x)) \rightarrow_{a.s.} 0,$$

for  $\varepsilon > 0$  small enough. Then the sequence of the MLEs is Hellinger consistent:

$$H(h \circ \hat{g}_n, h \circ g_0) \rightarrow_{a.s.} 0.$$

**PROOF.** For  $\varepsilon \in (0, 1)$  we have:

$$\begin{aligned} 0 &\geq \int_{\{h \circ g_0(x) \leq 1 - \varepsilon\}} \log(\varepsilon + h \circ g_0) dP_0 \geq \log(\varepsilon) P_0\{h \circ g_0(x) \leq 1 - \varepsilon\} > -\infty \\ 0 &\leq \int_{\{h \circ g_0(x) \geq 1\}} \log(\varepsilon + h \circ g_0) dP_0 \leq \int_{\{h \circ g_0(x) \geq 1\}} \log(2h \circ g_0) dP_0 \\ &\leq \int (h \log h) \circ g_0(x) dx + \log 2 < \infty. \end{aligned}$$

Thus the function  $\log(\varepsilon + h \circ g_0)$  is integrable with respect to probability measure  $P_0$ .



We can rearrange:

$$\begin{aligned}
 0 &\leq \mathbb{L}_n \hat{g}_n - \mathbb{L}_n g_0 = \int \log[h \circ \hat{g}_n] d\mathbb{P}_n - \int \log[h \circ g_0] d\mathbb{P}_n \\
 &\leq \int \log[\varepsilon + h \circ \hat{g}_n] d\mathbb{P}_n - \int \log[h \circ g_0] d\mathbb{P}_n \\
 (3.12) \quad &\leq \int \log[\varepsilon + h \circ \hat{g}_n] d(\mathbb{P}_n - P_0)
 \end{aligned}$$

$$(3.13) \quad + \int \log \left[ \frac{\varepsilon + h \circ \hat{g}_n}{\varepsilon + h \circ g_0} \right] dP_0$$

$$(3.14) \quad + \int \log[\varepsilon + h \circ g_0] dP_0 - \int \log[h \circ g_0] d\mathbb{P}_n.$$

The term (3.12) converges almost surely to zero by assumption.

For the term (3.13) we can apply the analogue of Lemma 1 from [Pal et al. \[2007\]](#):

$$II \equiv \int \log \left[ \frac{\varepsilon + h \circ \hat{g}_n}{\varepsilon + h \circ g_0} \right] dP_0 \leq 2 \int \sqrt{\frac{\varepsilon}{\varepsilon + h \circ g_0}} dP_0 - 2H^2(h \circ \hat{g}_n, h \circ g_0).$$

For the term (3.14), the SLLN implies that:

$$\begin{aligned}
 III &= \int \log[\varepsilon + h \circ g_0] dP_0 - \int \log[h \circ g_0] d\mathbb{P}_n \\
 &\rightarrow_{a.s.} \int \log[\varepsilon + h \circ g_0] dP_0 - \int \log[h \circ g_0] dP_0 = \int \log \left[ \frac{\varepsilon + h \circ g_0}{h \circ g_0} \right] dP_0.
 \end{aligned}$$

Thus we have:

$$\begin{aligned}
 0 &\leq \liminf(I + II + III) \\
 &\leq_{a.s.} -\limsup 2H^2(h \circ \hat{g}_n, h \circ g_0) + 2 \int \sqrt{\frac{\varepsilon}{\varepsilon + h \circ g_0}} dP_0 + \int \log \left[ \frac{\varepsilon + h \circ g_0}{h \circ g_0} \right] dP_0.
 \end{aligned}$$

This yields

$$\limsup H^2(h \circ \hat{g}_n, h \circ g_0) \leq_{a.s.} \int \sqrt{\frac{1}{1 + h \circ g_0/\varepsilon}} dP_0 + \frac{1}{2} \int \log \left[ \frac{\varepsilon + h \circ g_0}{h \circ g_0} \right] dP_0 \rightarrow 0$$

as  $\varepsilon \downarrow 0$  by monotone convergence.  $\square$

Next lemma allows us to obtain pointwise consistency once we proved Hellinger consistency.

**LEMMA 3.20.** *Consider that for a monotone model  $\mathcal{P}(h)$  a sequence of the MLEs  $\hat{g}_n$  is Hellinger consistent. Then the sequence  $\hat{g}_n$  is pointwise consistent. In other words  $\hat{g}_n(x) \rightarrow_{a.s.} g_0(x)$  for  $x \in \text{ri}(\text{dom } g_0)$  and convergence is uniform on compacta.*

PROOF. Let us denote by  $L_a^0$  and  $L_a^k$  the following sublevel sets:

$$\begin{aligned} L_a^0 &= \text{lev}_a g_0 \\ L_a^n &= \text{lev}_a \hat{g}_n. \end{aligned}$$

Consider  $\Omega_0$  such that  $\Pr[\Omega_0] = 1$  and  $H^2(h \circ \hat{g}_n^\omega, h \circ g_0) \rightarrow 0$  where  $\hat{g}_n^\omega$  is the MLE for  $\omega \in \Omega_0$ . For all  $\omega \in \Omega_0$  we have:

$$\begin{aligned} \int [\sqrt{h} \circ g_0 - \sqrt{h} \circ \hat{g}_n]^2 dx &\geq \int_{L_a^0 \setminus L_{a+\varepsilon}^n} [\sqrt{h} \circ g_0 - \sqrt{h} \circ \hat{g}_n]^2 dx \\ &\geq (\sqrt{h}(a) - \sqrt{h}(a + \varepsilon))^2 \mu(L_a^0 \setminus L_{a+\varepsilon}^n) \rightarrow 0, \end{aligned}$$

and by Lemma 4.2 we have:

$$\liminf \text{ri}(L_a^0 \cap L_{a+\varepsilon}^n) = \text{ri}(L_a^0).$$

Therefore  $\limsup \hat{g}_n(x) < a + \varepsilon$  for  $x \in \text{ri}(L_a^0)$ . Since  $a$  and  $\varepsilon$  are arbitrary we have  $\limsup \hat{g}_n \leq g_0$  on  $\text{ri}(\text{dom } g_0)$ .

On the other hand, we have:

$$\begin{aligned} \int [\sqrt{h} \circ g_0 - \sqrt{h} \circ \hat{g}_n]^2 dx &\geq \int_{L_{a-\varepsilon}^n \setminus L_a^0} [\sqrt{h} \circ g_0 - \sqrt{h} \circ \hat{g}_n]^2 dx \\ &\geq (\sqrt{h}(a - \varepsilon) - \sqrt{h}(a))^2 \mu(L_{a-\varepsilon}^n \setminus L_a^0) \rightarrow 0, \end{aligned}$$

and by Lemma 4.2 we have:

$$\limsup \text{cl}(L_{a-\varepsilon}^n \cup L_a^0) = \text{cl}(L_a^0).$$

Therefore  $\liminf \hat{g}_n(x) > a - \varepsilon$  for  $x$  such that  $g_0(x) \geq a$ . Since  $a$  and  $\varepsilon$  are arbitrary we have  $\liminf \hat{g}_n \geq g_0$  on  $\text{dom } g_0$ .

Thus  $\hat{g}_n \rightarrow g_0$  almost surely on  $\text{ri}(\text{dom } g_0)$ . By Theorem 10.8 Rockafellar [1970] convergence is uniform on compacta  $K \subset \text{ri}(\mathbb{R}_+^d)$ .  $\square$

We need a general property of the bracketing entropy numbers.

LEMMA 3.21. *Let  $\mathcal{A}$  be a class of sets in  $\mathbb{R}^d$  such that class  $\mathcal{A} \cap [-a, a]^d$  has finite bracketing entropy with respect to Lebesgue measure  $\lambda$  for any  $a$  large enough:*

$$\log N_{[]}(\varepsilon, \mathcal{A} \cap [-a, a]^d, L_1(\lambda)) < +\infty$$

*for every  $\varepsilon > 0$ . Then for any Lebesgue absolutely continuous probability measure  $P$  with bounded density we have that  $\mathcal{A}$  is Glivenko-Cantelli class:*

$$\|\mathbb{P}_n - P\|_{\mathcal{A}} \rightarrow_{a.s.} 0.$$

PROOF. Let  $C$  be an upper bound for the density of  $P$  and  $a$  be large so that for the set  $D \equiv [-a, a]^d$  we have  $P([-a, a]^d) > 1 - \varepsilon/2C$ . By assumption the class  $\mathcal{A} \cap D$  has a finite set of  $\varepsilon/2$ -brackets  $\{[L_i, U_i]\}$ . Then for any set  $A \in \mathcal{A}$  there exists index  $i$  such that:

$$L_i \subseteq A \cap D \subseteq U_i$$

Therefore:

$$L_i \subseteq A \subseteq U_i \cup D^c$$

and:

$$\begin{aligned} \|1\{U_i \cup D^c\} - 1\{L_i\}\|_{L_1(P)} &\leq \|1\{U_i\} - 1\{L_i\}\|_{L_1(P)} + \|1\{D^c\}\|_{L_1(P)} \\ &\leq C(\|1\{U_i\} - 1\{L_i\}\|_{L_1(\lambda)} + \|1\{D^c\}\|_{L_1(\lambda)}) \leq \varepsilon. \end{aligned}$$

Thus the set  $\{[L_i, U_i \cup D^c]\}$  is the set of  $\varepsilon$ -brackets for our class  $\mathcal{A}$  in  $L_1(P)$ . This implies that  $\mathcal{A}$  is a Glivenko-Cantelli class and the statement follows from Theorem 2.4.1 [van der Vaart and Wellner \[1996\]](#).  $\square$

To prove consistency for increasing models we begin with a general property of lower layer sets (see [Dudley \[1999\]](#), Chapter 8.3).

LEMMA 3.22. *Let  $\mathcal{LL}$  be the class of closed lower layer sets in  $\mathbb{R}_+^d$  and  $P$  be a Lebesgue absolutely continuous probability measure with bounded density. Then:*

$$\|\mathbb{P}_n - P\|_{\mathcal{LL}} \rightarrow_{a.s.} 0.$$

PROOF. By Theorem 8.3.2 [Dudley \[1999\]](#) we have

$$\log N_{[]}(\varepsilon, \mathcal{LL} \cap [0, 1]^d, L_1(\lambda)) < +\infty.$$

Since the class  $\mathcal{LL}$  is invariant under rescaling, the result follows from Lemma 3.21  $\square$

Note that Lemma 3.1 implies that if  $h \circ g$  belongs to an increasing model  $\mathcal{P}(h)$  then  $(\text{lev}_y g)^c$  is a lower layer set and has Lebesgue measure less or equal than  $1/h(y)$ . Let us denote by  $A_\delta$  the set  $\{|x| \leq \delta, x \in \mathbb{R}_+^d\}$ . Then by Lemma 3.1 part 3 we have:

$$(3.15) \quad (\text{lev}_y g)^c \subset A_{c/h(y)},$$

for  $c \equiv d!/d^d$ .

PROOF. (Theorem 2.16) By Lemma 3.19 it is enough to show that:

$$\int \log[\varepsilon + h \circ \hat{g}_n(x)] d(\mathbb{P}_n(x) - P_0(x)) \rightarrow_{a.s.} 0.$$

Indeed, applying Lemma 3.2 for the increasing transformation  $\log[\varepsilon + h(y)] - \log \varepsilon$  we obtain:

$$\begin{aligned} \int \log[\varepsilon + h \circ \hat{g}_n(x)] d(\mathbb{P}_n(x) - P_0(x)) &= \int_{-\infty}^{+\infty} \left[ \frac{h'(z)}{\varepsilon + h(z)} \right] (\mathbb{P}_n - P_0)(\text{lev}_z \hat{g}_n)^c dz \\ &\leq \|\mathbb{P}_n - P_0\|_{\mathcal{L}\mathcal{L}} \int_{-\infty}^M \left[ \frac{h'(z)}{\varepsilon + h(z)} \right] dz + \int_M^{+\infty} \left[ \frac{h'(z)}{\varepsilon + h(z)} \right] |\mathbb{P}_n - P_0|(\text{lev}_z \hat{g}_n)^c dz \\ &\leq \|\mathbb{P}_n - P_0\|_{\mathcal{L}\mathcal{L}} \log \left[ \frac{\varepsilon + h(M)}{\varepsilon} \right] + \int_M^{+\infty} \left[ \frac{h'(z)}{\varepsilon + h(z)} \right] (\mathbb{P}_n + P_0)(\text{lev}_z \hat{g}_n)^c dz. \end{aligned}$$

The first converges to zero almost surely by Lemma 3.22. For the second term we will use the inclusion 3.15:

$$\int_M^{+\infty} \left[ \frac{h'(z)}{\varepsilon + h(z)} \right] (\mathbb{P}_n + P_0)(\text{lev}_z \hat{g}_n)^c dz \leq \int_M^{+\infty} \left[ \frac{h'(z)}{\varepsilon + h(z)} \right] (\mathbb{P}_n + P_0) A_{c/h(z)} dz.$$

Now, we can apply Lemma 3.2 again for  $g_A(x) = h^{-1}(c/|\underline{x}|)$ . We have  $(\text{lev}_z g_A)^c = A_{c/h(z)}$  and therefore:

$$\begin{aligned} \int_M^{+\infty} \left[ \frac{h'(z)}{\varepsilon + h(z)} \right] (\mathbb{P}_n + P_0) A_{c/h(z)} dz &= \int_{A_{c/h(M)}} \log(\varepsilon + c/|\underline{x}|) d(\mathbb{P}_n + P_0) \\ &\leq \int_{A_{c/h(M)}} \log(2c/|\underline{x}|) d(\mathbb{P}_n + P_0), \end{aligned}$$

for  $M$  large enough. Assumption 1.5 and the SLLN imply that:

$$\int_{A_{c/h(M)}} \log(2c/|\underline{x}|) d(\mathbb{P}_n + P_0) \rightarrow_{a.s.} 2 \int_{A_{c/h(M)}} \log(2c/|\underline{x}|) dP_0.$$

Since  $M$  is arbitrary and  $A_{c/h(M)} \downarrow \{0\}$  as  $M \rightarrow +\infty$  the result follows.  $\square$

By Lemma 3.1 we have  $\text{ri}(\mathbb{R}_+^d) \subseteq \text{dom } g_0$ . Thus Theorem 2.16 and Lemma 3.20 imply Theorem 2.17.

**THEOREM 3.23.** *For an increasing model  $\mathcal{P}(h)$  and a true density  $h \circ g$  which satisfies the assumptions 1.4, 1.5 and 1.6 the sequence of MLEs  $\hat{g}_n$  is pointwise consistent. That is  $\hat{g}_n(x) \rightarrow_{a.s.} g_0(x)$  for  $x \in \text{ri}(\mathbb{R}_+^d)$  and convergence is uniform on compacta.*

Finally, we prove consistency for decreasing models. We need a general property of convex sets.

LEMMA 3.24. *Let  $\mathcal{A}$  be the class of closed convex sets  $A$  in  $\mathbb{R}^d$  and  $P$  be a Lebesgue absolutely continuous probability measure with bounded density. Then:*

$$\|\mathbb{P}_n - P\|_{\mathcal{A}} \rightarrow_{a.s.} 0.$$

PROOF. Let  $D$  be a convex compact set. By Theorem 8.4.2 [Dudley \[1999\]](#) the class  $\mathcal{A} \cap D$  has a finite set of  $\varepsilon$ -brackets. Since the class  $\mathcal{A}$  is invariant under rescaling, the result follows from Lemma [3.21](#)  $\square$

LEMMA 3.25. *For a decreasing model  $\mathcal{P}(h)$  the sequence of MLEs  $\hat{g}_n$  is almost surely uniformly bounded below.*

PROOF. We will apply Lemma [3.17](#) to the sequences  $\hat{g}_n$  and  $\{n\}$ . By the SLLN and Lemma [3.9](#) we have:

$$\mathbb{L}_n \hat{g}_n \geq \mathbb{L}_n g_0 \rightarrow_{a.s.} \int [h \log h] \circ g_0 dx > -\infty.$$

Therefore the sequence  $\{\mathbb{L}_n \hat{g}_n\}$  is bounded away from  $-\infty$ , and the first condition of Lemma [3.17](#) is true.

Choose some  $a \in (0, d/n_d)$ . Then for any set  $S$  such that  $\mu[S] = \rho \equiv a/h(\min g_0)$  where  $\min g_0$  is attained by Lemma [3.6](#) we have:

$$P[S] = \int_S h \circ g_0 dx \leq \mu[S] h(\min g_0) = a < d/n_d.$$

Now, let  $A_n = \text{lev}_{a_n} \hat{g}_n$  be such sets that  $\mu[A_n] = \rho$ . Then, by Lemma [3.24](#) we have:

$$|\mathbb{P}_n[A_n] - P[A_n]| \leq \|\mathbb{P}_n - P\|_{\mathcal{A}} \rightarrow_{a.s.} 0,$$

which implies that  $\mathbb{P}_n[A_n] < d/n_d$  almost surely for  $n$  large enough. Therefore, the second condition of Lemma [3.17](#) is true and it is applicable to the sequence  $\hat{g}_n$  almost surely.  $\square$

PROOF. (Theorem [2.18](#)) By Lemma [3.9](#) and Lemma [3.19](#) it is enough to show that:

$$\int \log[\varepsilon + h \circ \hat{g}_n(x)] d(\mathbb{P}_n(x) - P_0(x)) \rightarrow_{a.s.} 0.$$

By Lemma [3.25](#) we have  $\inf \hat{g}_n \geq A$  for some  $A > h_\infty$ . Therefore, by Lemma [3.7](#) applied to the decreasing transformation  $\log[\varepsilon + h(y)] - \log \varepsilon$

it follows:

$$\begin{aligned}
\int \log[\varepsilon + h \circ \hat{g}_n(x)] d(\mathbb{P}_n(x) - P_0(x)) &= \int_A^{+\infty} \left[ \frac{-h'(z)}{\varepsilon + h(z)} \right] (\mathbb{P}_n - P_0)(\text{lev}_z \hat{g}_n) dz \\
&\leq \|\mathbb{P}_n - P_0\|_{\mathcal{A}} \int_A^{+\infty} \left[ \frac{-h'(z)}{\varepsilon + h(z)} \right] dz \\
&= \|\mathbb{P}_n - P_0\|_{\mathcal{A}} \log \left[ \frac{\varepsilon + h(A)}{\varepsilon} \right] \rightarrow_{a.s.} 0,
\end{aligned}$$

where the last limit follows from Lemma 3.24.  $\square$

PROOF. (Theorem 2.19) By Lemma 3.20  $\hat{g}_n \rightarrow g_0$  almost surely on  $\text{ri}(\text{dom } g_0)$ . Functions  $g_0$  and  $g_0^*$  differ only on the boundary  $\partial \text{dom } g_0$  which has Lebesgue measure zero by Lemma 4.1. Since observations  $X_i \in \text{ri}(\text{dom } g_0)$  almost surely we have  $\hat{g}_n = +\infty$  on  $\partial \text{dom } g_0$  and thus  $\hat{g}_n \rightarrow g_0^*$ .

Now, we assume that  $\text{dom } g_0 = \mathbb{R}^d$ . By Lemma 3.6 function  $g_0$  has bounded sublevel sets and therefore there exists  $x_0$  where  $g_0$  attains its minimum  $m$ . Since  $h \circ g_0$  is density we have  $h(m) > 0$  and by Lemma 3.8 we have  $h(m) < \infty$ . Fix  $\varepsilon > 0$  such that  $h(m) > 3\varepsilon$  and consider  $a$  such that  $h(a) < \varepsilon$ . The set  $A = \text{lev}_a g_0$  is bounded and by continuity  $g_0 = a$  on  $\partial A$ . Choose  $\delta > 0$  such that  $h(a - \delta) < 2\varepsilon < h(m + \delta)$  and:

$$\sup_{x \in [m, a + \delta]} |h(x) - h(x - \delta)| \leq \varepsilon.$$

The closure  $\bar{A}$  is compact and thus for  $n$  large enough we have with probability one:

$$\sup_{\bar{A}} |\hat{g}_n - g_0| < \delta,$$

which implies:

$$\sup_{\bar{A}} |h \circ \hat{g}_n - h \circ g_0| < \varepsilon,$$

since the range of values of  $g_0$  on  $\bar{A}$  is  $[m, a]$ . The set  $\partial A$  is compact and therefore  $\hat{g}_n$  attains its minimum  $m_n$  on this set at some point  $x_n$ . By construction:

$$m_n = \hat{g}_n(x_n) > g_0(x_n) - \delta = a - \delta > m + \delta = g_0(x_0) + \delta > \hat{g}_n(x_0) = m$$

We have  $x_0 \in A \cap \text{lev}_{a-\delta} \hat{g}_n$  and  $\hat{g}_n \geq m_n > a - \delta$  on  $\partial A$ . Thus, by convexity we have  $\text{lev}_{a-\delta} \hat{g}_n \subset A$  and for  $x \notin \bar{A}$  we have:

$$|h \circ \hat{g}_n(x) - h \circ g_0(x)| \leq h \circ \hat{g}_n(x) + h \circ g_0(x) < h(a - \delta) + h(a) < 3\varepsilon.$$

This shows that for any  $\varepsilon > 0$  small enough we will have:

$$\|h \circ \hat{g}_n - h \circ g_0\|_\infty < 3\varepsilon$$

with probability one as  $n \rightarrow \infty$ . This concludes the proof.  $\square$

**3.5. Proofs for Lower Bound Results.** We will use the following lemma for computing the Hellinger distance between a function and its local deformation:

**LEMMA 3.26.** *Let  $\{g_\varepsilon\}$  be a local deformation of the function  $g : \mathbb{R}^d \rightarrow \mathbb{R}$  at the point  $x_0$ , such that  $g$  is continuous at  $x_0$ , and let the function  $h : \mathbb{R} \rightarrow \mathbb{R}$  be continuously differentiable at the point  $g(x_0)$ . Then for any  $r > 0$ :*

$$(3.16) \quad \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^d} |g_\varepsilon(x) - g(x)|^r dx = 0,$$

$$(3.17) \quad \lim_{\varepsilon \rightarrow 0} \frac{\int_{\mathbb{R}^d} |h \circ g_\varepsilon(x) - h \circ g(x)|^r dx}{\int_{\mathbb{R}^d} |g_\varepsilon(x) - g(x)|^r dx} = |h' \circ g(x_0)|^r.$$

**PROOF.** Since  $\{g_\varepsilon\}$  is a local deformation, for  $\varepsilon > 0$  small enough we have:

$$\begin{aligned} \int_{\mathbb{R}^d} |h \circ g_\varepsilon(x) - h \circ g(x)|^r dx &= \int_{B(x_0; r_\varepsilon)} |h \circ g_\varepsilon(x) - h \circ g(x)|^r dx, \\ \int_{\mathbb{R}^d} |g_\varepsilon(x) - g(x)|^r dx &= \int_{B(x_0; r_\varepsilon)} |g_\varepsilon(x) - g(x)|^r dx. \end{aligned}$$

Then:

$$\int_{B(x_0; r_\varepsilon)} |g_\varepsilon - g|^r dx \leq \text{ess sup } |g_\varepsilon - g|^r \mu[B(x_0; r_\varepsilon)]$$

implies (3.16).

Let us define a sequence  $\{a_\varepsilon\}$ :

$$a_\varepsilon \equiv \text{ess sup } |g_\varepsilon - g| + \sup_{x \in B(x_0; r_\varepsilon)} |g(x) - g(x_0)|.$$

For  $x \in B(x_0; r_\varepsilon)$  and  $y \in [g_\varepsilon(x), g(x)]$  we have a.e.:

$$|y - g(x_0)| \leq |g_\varepsilon(x) - g(x)| + |g(x) - g(x_0)| \leq a_\varepsilon.$$

Using the mean value theorem we obtain:

$$\begin{aligned} \int_{\mathbb{R}^d} |h \circ g_\varepsilon(x) - h \circ g(x)|^r dx &= \int_{\mathbb{R}^d} |h'(y)|^r |g_\varepsilon(x) - g(x)|^r dx \\ \inf_{y \in B(g(x_0); a_\varepsilon)} |h'(y)|^r &\leq \frac{\int_{\mathbb{R}^d} |h \circ g_\varepsilon(x) - h \circ g(x)|^r dx}{\int_{\mathbb{R}^d} |g_\varepsilon(x) - g(x)|^r dx} \leq \sup_{y \in B(g(x_0); a_\varepsilon)} |h'(y)|^r \end{aligned}$$

Since  $h'$  is continuous at  $g(x_0)$ , to prove (3.17) it is enough to show that  $a_\varepsilon \rightarrow 0$ . By assumption we have:

$$\lim_{\varepsilon \rightarrow 0} \operatorname{ess\,sup} |g_\varepsilon - g| = 0.$$

Since  $g$  is continuous at  $x_0$  and  $r_\varepsilon \rightarrow 0$  we have:

$$\lim_{\varepsilon \rightarrow 0} \sup_{x \in B(x_0; r_\varepsilon)} |g(x) - g(x_0)| = 0.$$

Thus  $a_\varepsilon \rightarrow 0$ , which proves (3.17).  $\square$

In order to apply Corollary 2.22 we need to construct deformations so that they still belong to the class  $\mathcal{G}$ . The following lemma provides a technique for constructing such deformations.

LEMMA 3.27. *Let  $\{g_\varepsilon\}$  be a local deformation of the function  $g : \mathbb{R}^d \rightarrow \mathbb{R}$  at the point  $x_0$ , such that  $g$  is continuous at  $x_0$ , and let the function  $h : \mathbb{R} \rightarrow \mathbb{R}$  be continuously differentiable at the point  $g(x_0)$  so that  $h' \circ g(x_0) \neq 0$ . Then for any fixed  $\delta > 0$  small enough, the deformation  $g_{\theta, \delta} = \theta g_\delta + (1 - \theta)g$  and any  $r > 0$  we have:*

$$(3.18) \quad \limsup_{\theta \rightarrow 0} \theta^{-r} \int_{\mathbb{R}^d} |h \circ g_{\theta, \delta}(x) - h \circ g(x)|^r dx < \infty,$$

$$(3.19) \quad \liminf_{\theta \rightarrow 0} \theta^{-r} \int_{\mathbb{R}^d} |h \circ g_{\theta, \delta}(x) - h \circ g(x)|^r dx > 0.$$

Note that  $g_{\theta, \delta}$  is not a local deformation.

PROOF. The statement follows from the argument for Lemma 3.26. For a fixed  $\theta$  the family  $\{g_{\theta, \delta}\}$  is a local deformation. Thus for  $a_{\theta, \varepsilon}$  defined by:

$$a_{\theta, \varepsilon} \equiv \operatorname{ess\,sup} |g_{\theta, \varepsilon} - g| + \sup_{x \in B(x_0; r_\varepsilon)} |g(x) - g(x_0)|,$$

it follows that

$$\begin{aligned} \frac{\int_{\mathbb{R}^d} |h \circ g_{\theta, \varepsilon}(x) - h \circ g(x)|^r dx}{\int_{\mathbb{R}^d} |g_{\theta, \varepsilon}(x) - g(x)|^r dx} &\leq \sup_{y \in B(g(x_0); a_{\theta, \varepsilon})} |h'(y)|^r, \\ \frac{\int_{\mathbb{R}^d} |h \circ g_{\theta, \varepsilon}(x) - h \circ g(x)|^r dx}{\int_{\mathbb{R}^d} |g_{\theta, \varepsilon}(x) - g(x)|^r dx} &\geq \inf_{y \in B(g(x_0); a_{\theta, \varepsilon})} |h'(y)|^r. \end{aligned}$$

For  $|\theta| < 1$  we have:

$$|g_{\theta, \varepsilon} - g| = |\theta| |g_\varepsilon - g|$$



and therefore  $a_{\theta,\delta} \leq a_\delta$ . Since  $a_\varepsilon \rightarrow 0$  and  $h$  is continuously differentiable for all  $\delta > 0$  small enough we have:

$$\begin{aligned} \sup_{y \in B(g(x_0); a_{\theta,\delta})} |h'(y)|^r &\leq \sup_{y \in B(g(x_0); a_\delta)} |h'(y)|^r < \infty, \\ \inf_{y \in B(g(x_0); a_{\theta,\delta})} |h'(y)|^r &\geq \inf_{y \in B(g(x_0); a_\delta)} |h'(y)|^r > 0. \end{aligned}$$

Thus for all  $\theta$  we obtain:


$$\begin{aligned} \theta^{-r} \int_{\mathbb{R}^d} |h \circ g_{\theta,\delta}(x) - h \circ g(x)|^r dx \\ \leq \sup_{y \in B(g(x_0); a_\delta)} |h'(y)|^r \int_{\mathbb{R}^d} |g_\delta(x) - g(x)|^r dx < \infty, \\ \theta^{-r} \int_{\mathbb{R}^d} |h \circ g_{\theta,\delta}(x) - h \circ g(x)|^r dx \\ \geq \inf_{y \in B(g(x_0); a_\delta)} |h'(y)|^r \int_{\mathbb{R}^d} |g_\delta(x) - g(x)|^r dx > 0 \end{aligned}$$

which proves the lemma.  $\square$

PROOF. (Theorem 2.24). Our statement is not trivial only if the curvature  $\text{curv}_{x_0} g > 0$  or equivalently there exists positive definite  $d \times d$  matrix  $G$  such that the function  $g$  is locally  $G$ -strongly convex. Then by Lemma 4.17 this means that there exists a convex function  $q$  such that in some neighborhood  $O(x_0)$  of  $x_0$  we have:

$$(3.20) \quad g(x) = \frac{1}{2}(x - x_0)^T G(x - x_0) + q(x).$$

The plan of the proof is the following: we introduce families of functions  $\{D_\varepsilon(g; x_0, v)\}$  and  $\{D_\varepsilon^*(g; x_0)\}$  and prove that these families are local deformations. Using these deformations as building blocks we construct two types of deformations:  $\{h \circ g_\varepsilon^+\}$  and  $\{h \circ g_\varepsilon^-\}$  of the density  $h \circ g$  which belong to  $\mathcal{P}(h)$ . These deformations represent positive and negative changes in the value of the function  $g$  at the point  $x_0$ . After that we approximate the Hellinger distances using Lemma 3.26. Finally, applying Corollary 2.22 we obtain lower bounds which depend on  $G$ . We finish the proof by taking the supremum of the obtained lower bounds over all  $G \in \mathcal{SC}(g; x_0)$ . Under the mild assumption of strong convexity of the function  $g$  both deformations give the same rate and structure of the constant  $C(d)$ . However, it is possible to obtain a larger constant  $C(d)$  for the negative deformation if we assume that  $g$  is twice differentiable. Note that by the definition of  $\mathcal{P}(h)$  the function  $g$  is a closed proper convex function.

FIG 1. Example of the deformation  $D_\varepsilon(g; x_0, v_0)$ .

Let us define a function  $D_\varepsilon(g; x_0, v_0)$  for a given  $\varepsilon > 0$ ,  $x_0 \in \text{dom } g$  and  $v_0 \in \partial g(x_0)$  as:

$$D_\varepsilon(g; x_0, v_0)(x) = \max(g(x), l_0(x) + \varepsilon),$$

where  $l_0(x) = \langle v_0, x - x_0 \rangle + g(x_0)$  is a support plane to  $g$  at  $x_0$ . Since  $l_0 + \varepsilon$  is a support plane to  $g + \varepsilon$  we have:

$$g \leq D_\varepsilon(g; x_0, v_0) \leq g + \varepsilon,$$

and thus  $\text{dom } D_\varepsilon(g; x_0, v_0) = \text{dom } g$ . As a maximum of two closed convex functions  $D_\varepsilon(g; x_0, v_0)$  is a closed convex function. For a given  $x_1$  we have  $D_\varepsilon(g; x_0, v_0)(x_1) = g(x_1)$  if and only if:


$$(3.21) \quad g(x_1) - \varepsilon \geq \langle v_0, x_1 - x_0 \rangle + h(x_0).$$

We also define a function  $D_\varepsilon^*(g; x_0)$  for a given  $\varepsilon > 0$  and  $x_0 \in \text{dom } g$  as a maximal convex minorant (Appendix 4.1) of the function  $\tilde{g}_\varepsilon$  defined as:

$$\tilde{g}_\varepsilon(x) = \begin{cases} g(x), & x \neq x_0 \\ g(x_0) - \varepsilon, & x = x_0. \end{cases}$$

Both functions  $D_\varepsilon(g; x_0, v_0)$  and  $D_\varepsilon^*(g; x_0)$  are convex by construction, and, as the next lemma shows, have similar properties. However, the argument for  $D_\varepsilon^*(g; x_0)$  is more complicated.

**LEMMA 3.28.** *Let  $g$  be a closed proper convex function,  $g^*$  be its convex conjugate and  $x_0 \in \text{ri}(\text{dom } g)$ . Then:*

FIG 2. Example of the deformation  $D_\varepsilon^*(g; x_0)$ .

1.  $D_\varepsilon^*(g; x_0)$  is a closed proper convex function such that:

$$g - \varepsilon \leq D_\varepsilon^*(g; x_0) \leq g$$

and  $\text{dom } D_\varepsilon^*(g; x_0) = \text{dom } g$ .

2. For a given  $x_1 \in \text{ri}(\text{dom } g)$  we have  $D_\varepsilon^*(g; x_0)(x_1) = g(x_1)$  if and only if there exists  $v \in \partial g(x_1)$  such that:

$$(3.22) \quad g(x_1) + \varepsilon \leq \langle v, x_1 - x_0 \rangle + g(x_0).$$

3. If  $y_0 \in \partial g(x_0)$  then  $x_0 \in \partial g^*(y_0)$  and:

$$D_\varepsilon(g; x_0, y_0) = (D_\varepsilon^*(g^*; y_0))^*.$$

PROOF. Obviously,  $\tilde{g}_\varepsilon \geq g - \varepsilon$ . Since  $g - \varepsilon$  is a closed proper convex function it is equal to the supremum of all linear functions  $l$  such that  $l \leq g - \varepsilon$ . Thus  $g - \varepsilon \leq D_\varepsilon^*(g; x_0)$ , which implies that  $D_\varepsilon^*(g; x_0)$  is a proper convex function and  $\text{dom } D_\varepsilon^*(g; x_0) \subseteq \text{dom}(g - \varepsilon) = \text{dom } g$ . By Lemma 4.10 we have  $D_\varepsilon^*(g; x_0) \leq g$ , therefore  $\text{dom } g \subseteq \text{dom } D_\varepsilon^*(g; x_0)$  which proves 1.

If  $v \in \partial g(x_1)$  then  $l_v(x) = \langle v, x - x_1 \rangle + g(x_1)$  is a support plane to  $g(x)$  and  $l_v \leq g$ . If inequality (3.22) is true then  $l_v(x)$  is majorized by  $\tilde{g}_\varepsilon$  and we have:

$$D_\varepsilon(g; x_0)(x_1) \leq g(x_1) = l_v(x_1) \leq D_\varepsilon(g; x_0)(x_1).$$

On the other hand, by 1 we have  $x_1 \in \text{ri}(\text{dom } D_\varepsilon(g; x_0))$ , hence there exists  $v \in \partial D_\varepsilon(g; x_0)(x_1)$  and:

$$\begin{aligned} g(x) \geq \tilde{g}_\varepsilon(x) \geq D_\varepsilon(g; x_0)(x) &\geq \langle v, x_0 - x_1 \rangle + D_\varepsilon(g; x_0)(x_1) \\ &= \langle v, x_0 - x_1 \rangle + g(x_1). \end{aligned}$$

Therefore  $v \in \partial g(x_1)$ . In particular:

$$\begin{aligned} \tilde{g}_\varepsilon(x_0) &= g(x_0) - \varepsilon \geq D_\varepsilon(g; x_0)(x_0) \\ &\geq \langle v, x_0 - x_1 \rangle + D_\varepsilon(g; x_0)(x_1) = \langle v, x_0 - x_1 \rangle + g(x_1), \end{aligned}$$

which proves 2.

We can represent  $D_\varepsilon^*(g^*; x_0)$  as the maximal convex minorant of  $g$  defined by:

$$g = \min(g, g(x_0) - \varepsilon + \delta(\cdot | x_0)).$$

For  $x \in \text{dom } g$  by Lemma 4.10,  $g^*(y_0) + g(x_0) = \langle y_0, x_0 \rangle$ . Thus

$$(g(x_0) - \varepsilon + \delta(\cdot | x_0))^*(y) = \langle x_0, y \rangle - g(x_0) + \varepsilon = \langle x_0, y - v \rangle + \varepsilon$$

for some  $v \in \partial g(x_0)$ . By Lemma 4.7 we have:

$$D_\varepsilon^*(g^*; x_0)^* = \max(g^*, l_0^*), \quad l_0^*(y) \equiv \langle x_0, y - v \rangle + \varepsilon$$

which concludes the proof the lemma.  $\square$

Since the domain of the quadratic part of the equation (3.20) is  $\mathbb{R}^d$ , by Lemma 4.11 we have that for any  $x \in \text{dom } g$  and  $v \in \partial g(x)$  there exists  $w \in \partial q(x)$  such that:

$$(3.23) \quad v = G(x - x_0) + w.$$

Therefore for the point  $x_1$  in the neighborhood  $O(x_0)$  where the decomposition (3.20) is true, condition (3.21) is equivalent to:

$$\frac{1}{2}(x_1 - x_0)^T G(x_1 - x_0) + q(x_1) - \varepsilon \geq \langle w_0, x_1 - x_0 \rangle + q(x_0),$$

where  $w_0$  corresponds to  $y_0$  in the decomposition (3.23). Since  $\langle w_0, x_1 - x_0 \rangle + q(x_0)$  is a support plane to  $q(x)$ , the inequality (3.21) is satisfied if

$$\frac{1}{2}(x_1 - x_0)^T G(x_1 - x_0) \geq \varepsilon,$$

which is the complement of an open ellipsoid  $B_G(x_0, \sqrt{2\varepsilon})$  defined by  $G$  with the center at  $x_0$ . For  $\varepsilon$  small enough this ellipsoid will belong to the neighborhood  $O(x_0)$ . Since  $|D_\varepsilon(g; x_0, y_0) - g| \leq \varepsilon$  this proves that the family  $D_\varepsilon(g; x_0, y_0)$  is a local deformation.

In the same way, the condition (3.22) is equivalent to:

$$\frac{1}{2}(x_1 - x_0)^T G(x_1 - x_0) + q(x_1) + \varepsilon \leq \langle G(x_1 - x_0) + w_1, x_1 - x_0 \rangle + q(x_0)$$

or:

$$\frac{1}{2}(x_1 - x_0)^T G(x_1 - x_0) + q(x_0) - \varepsilon \geq \langle w_1, x_0 - x_1 \rangle + q(x_1),$$

which is satisfied if we have:

$$\frac{1}{2}(x_1 - x_0)^T G(x_1 - x_0) \geq \varepsilon.$$

Since  $|D_\varepsilon^*(g; x_0) - g| \leq \varepsilon$  this proves that the family  $D_\varepsilon^*(g; x_0)$  is also a local deformation. Thus we proved:

LEMMA 3.29. *Let  $g$  be a closed proper convex function locally  $G$ -strongly convex at some  $x_0 \in \text{ri dom } g$  and  $y_0 \in \partial g(x_0)$ . Then the families  $D_\varepsilon(g; x_0, y_0)$  and  $D_\varepsilon^*(g; x_0)$  are local deformations for all  $\varepsilon > 0$  small enough. Moreover, the condition:*

$$\frac{1}{2}(x - x_0)^T G(x - x_0) \geq \varepsilon$$

*implies*

$$D_\varepsilon(g; x_0, y_0)(x) = D_\varepsilon^*(g; x_0)(x) = g(x).$$

*Or equivalently  $\text{supp}[D_\varepsilon(g; x_0, y_0) - g]$  and  $\text{supp}[D_\varepsilon^*(g; x_0) - g]$  are subsets of  $B_G(x_0, \sqrt{2\varepsilon})$ .*

For  $r > 0$  small enough  $h' \circ g(x)$  is positive and the decomposition 3.20 is true on  $B(x_0; r)$ . Let us fix some  $y_0 \in \partial g(x_0)$ , some  $x_1 \in B(x_0; r)$  such that  $x_1 \neq x_0$ , and some  $y_1 \in \partial g(x_1)$ . We fix  $\delta$  such that equation (3.18) of Lemma 3.27 is true for the transformation  $\sqrt{h}$  and  $r = 2$  and also  $x_0 \notin B_G(x_1; \sqrt{2\delta})$ . Then by Lemma 3.29 for all  $\varepsilon > 0$  small enough, the support sets  $\text{supp}[D_\varepsilon(g; x_0, y_0) - g]$  and  $\text{supp}[D_\varepsilon^*(g; x_0) - g]$  do not intersect; i.e. these two deformations do not interfere.

Now, we can prove Theorem 2.24. The argument below is identical for  $g_\varepsilon^+$  and  $g_\varepsilon^-$ , so we will give the proof only for  $g_\varepsilon^+$ . We define deformations  $g_\varepsilon^+$  and  $g_\varepsilon^-$  by means of the following Lemma:

LEMMA 3.30. *For all  $\varepsilon > 0$  small enough there exist  $\theta_\varepsilon^+, \theta_\varepsilon^- \in (0, 1)$  such that the functions  $g_\varepsilon^+$  and  $g_\varepsilon^-$  defined by:*

$$\begin{aligned} g_\varepsilon^+ &= (1 - \theta_\varepsilon^+)D_\varepsilon(g; x_0, v_0) + \theta_\varepsilon^+D_\delta^*(g; x_1) \\ g_\varepsilon^- &= (1 - \theta_\varepsilon^-)D_\varepsilon^*(g; x_0) + \theta_\varepsilon^-D_\delta(g; x_1; v_1) \end{aligned}$$

*belong to  $\mathcal{P}(h)$ .*

PROOF. By dominated convergence, the function  $F(\theta)$  defined by:

$$F(\theta) = \int h \circ ((1 - \theta)D_\varepsilon(g; x_0, v_0) + \theta D_\delta^*(g; x_1)) dx$$

is continuous. We have:

$$\begin{aligned} F(0) &= \int h \circ D_\varepsilon(g; x_0, v_0) dx > \int h \circ g dx = 1, \\ F(1) &= \int h \circ D_\delta^*(g; x_1) dx < \int h \circ g dx = 1. \end{aligned}$$

Therefore there exists  $\theta_\varepsilon^+ \in (0, 1)$  such that  $F(\theta_\varepsilon^+) = 1$ .  $\square$

Next, we will show that  $\theta_\varepsilon^+$  goes to zero fast enough so that  $g_\varepsilon^+$  is very close to  $D_\varepsilon(g; x_0, v_0)$ . Since supports do not intersect we have:

$$\begin{aligned} 0 &= \int (h \circ g_\varepsilon^+ - h \circ g) dx = \int (h \circ ((1 - \theta_\varepsilon^+)D_\varepsilon(g; x_0, v_0) + \theta_\varepsilon^+g) - h \circ g) dx \\ &\quad - \int (h \circ g - h \circ ((1 - \theta_\varepsilon^+)g + \theta_\varepsilon^+D_\delta^*(g; x_1))) dx, \end{aligned}$$

where both integrals have the same sign. For the first integral by Lemma 3.26 we have:

$$\begin{aligned} &\int |h \circ ((1 - \theta_\varepsilon^+)D_\varepsilon(g; x_0, v_0) + \theta_\varepsilon^+g) - h \circ g| dx \\ &\leq \int |h \circ D_\varepsilon(g; x_0, v_0) - h \circ g| dx \\ &\asymp \int (g - D_\varepsilon(g; x_0, v_0)) dx \leq \varepsilon \mu[B_G(x_0; \sqrt{2\varepsilon})]. \end{aligned}$$

The second integral is monotone in  $\theta_\varepsilon^+$ , and by Lemma 3.27 we have:

$$\int (h \circ g - h \circ ((1 - \theta_\varepsilon^+)g + \theta_\varepsilon^+D_\delta^*(g; x_1))) dx \asymp \theta_\varepsilon^+.$$

Thus we have  $\theta_\varepsilon^+ = O(\varepsilon^{1+d/2})$  and:

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} (g_\varepsilon^+(x_0) - g(x_0)) = \lim_{\varepsilon \rightarrow 0} (1 - \theta_\varepsilon^+) = 1.$$

For Hellinger distance we have:

$$\begin{aligned} H(h \circ g_\varepsilon^m, h \circ g) &= H(h \circ ((1 - \theta_\varepsilon^+) D_\varepsilon(g; x_0, v_0) + \theta_\varepsilon^+ g), h \circ g) \\ &\quad + H(h \circ ((1 - \theta_\varepsilon^+) g + \theta_\varepsilon^+ D_\delta^*(g; x_1)), h \circ g). \end{aligned}$$

Now, we can apply Lemma 3.26

$$\begin{aligned} H^2(h \circ ((1 - \theta_\varepsilon^+) D_\varepsilon(g; x_0, v_0) + \theta_\varepsilon^+ g), h \circ g) &\leq H^2(h \circ D_\varepsilon(g; x_0, v_0), h \circ g), \\ \lim_{\varepsilon \rightarrow 0} \frac{H^2(h \circ D_\varepsilon(g; x_0, v_0), h \circ g)}{\int (D_\varepsilon(g; x_0, v_0) - g)^2 dx} &= \frac{h' \circ g(x_0)^2}{4h \circ g(x_0)}, \end{aligned}$$

and

$$\int (D_\varepsilon(g; x_0, v_0) - g)^2 dx \leq \varepsilon^2 \mu[B_G(x_0; \sqrt{2\varepsilon})] = \varepsilon^{2+d/2} \frac{2^{d/2} \mu[S(0, 1)]}{\sqrt{\det G}}.$$

This yields

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \varepsilon^{-\frac{d+4}{4}} H(h \circ ((1 - \theta_\varepsilon^+) D_\varepsilon(g; x_0, v_0) + \theta_\varepsilon^+ g), h \circ g) \\ \leq C(d) \left( \frac{h' \circ g(x_0)^4}{h \circ g(x_0)^2 \det G} \right)^{1/4} \end{aligned}$$

where  $S(0, 1)$  is  $d$ -dimensional sphere of radius 1.

For the second part by Lemma 3.27 we obtain:

$$\limsup_{\varepsilon \rightarrow 0} (\theta_\varepsilon^+)^{-2} H^2(h \circ ((1 - \theta_\varepsilon^+) g + \theta_\varepsilon^+ D_\delta^*(g; x_1)), h \circ g) < \infty,$$

and

$$H(h \circ ((1 - \theta_\varepsilon^+) g + \theta_\varepsilon^+ D_\delta^*(g; x_1)), h \circ g) = O(\varepsilon^{\frac{d+2}{2}}).$$

Thus:

$$\limsup_{\varepsilon \rightarrow 0} \varepsilon^{-\frac{d+4}{4}} H(h \circ g_\varepsilon^+, h \circ g) \leq C(d) \left( \frac{h' \circ g(x_0)^4}{h \circ g(x_0)^2 \det G} \right)^{1/4}.$$

Finally, we apply Corollary 2.22:

$$\liminf_{n \rightarrow \infty} n^{\frac{2}{d+4}} R_1(n; T, \{g, g_n\}) \geq C(d) \left[ \frac{h \circ g(x_0)^2 \det G}{h' \circ g(x_0)^4} \right]^{\frac{1}{d+4}}.$$

Taking the supremum over all  $G \in \mathcal{SC}(g; x_0)$  we obtain the statement of the theorem.  $\square$

### 3.6. Mode estimation.

PROOF. (Theorem 2.27). The proof is similar to the proof for a point estimation lower bounds. The deformation we will construct will resemble  $g_\varepsilon^-$ .

Our statement is not trivial only if the curvature  $\text{curv}_{x_0} g > 0$  or equivalently there exists such positive definite  $d \times d$  matrix  $G$  so that the function  $g$  is locally  $G$ -strongly convex. For  $a > 0$  small enough  $h' \circ g(x)$  is positive and decomposition 3.20 is true on  $B(x_0; a)$ . Let us fix some  $y_0 \in \partial g(x_0)$ , some  $x_1 \in B(x_0; a)$  such that  $x_1 \neq x_0$  and some  $y_1 \in \partial g(x_1)$ . We fix  $\delta$  such that equation (3.18) of Lemma 3.27 is true for the transformation  $\sqrt{h}$  and  $r = 2$  and also  $x_0 \notin \overline{B_G(x_1; \sqrt{2\delta})}$ .

Let us consider the deformation  $D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u)$  where  $u \in \mathbb{R}^d$  is an arbitrary fixed vector in  $\mathbb{R}^d$  with  $\|u\| = 1$  and

$$\xi(\varepsilon) = g(x_0) - g(x_0 + \varepsilon u) + \varepsilon^{\gamma+1}.$$

Since the value of  $D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u)$  at any point  $x$  is a convex combination of  $g(y)$  for some  $y$ ,  $g(x) \geq g(x_0)$  and

$$D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u)(x_0 + \varepsilon u) = g(x_0) + \varepsilon^{\gamma+1}$$

the global minimum of  $D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u)$  is  $x_0 + \varepsilon u$ . By Lemma 3.29 for all  $\varepsilon > 0$  small enough we have

$$\text{supp}[D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) - g] \subseteq B_G(x_0 + \varepsilon u, \sqrt{2\xi(\varepsilon)}).$$

Since, by assumption

$$\xi(\varepsilon) \leq L\varepsilon^\gamma + \varepsilon^{\gamma+1}$$

the support of  $\text{supp}[D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) - g]$  converges to a point  $x_0$  and thus does not intersect  $\text{supp}[D_\varepsilon(g; x_1, y_1) - g]$  for  $\varepsilon$  small enough i.e. these two deformations do not interfere.

The same argument as in Lemma 3.30 shows that there exists  $\theta_\varepsilon^m \in (0, 1)$  such that the deformation  $g_\varepsilon^m$  defined as:

$$g_\varepsilon^m = (1 - \theta_\varepsilon^m)D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) + \theta_\varepsilon^m D_\delta(g; x_1, y_1)$$

belongs to  $\mathcal{P}(h)$ . Also  $g_\varepsilon^m \geq D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u)$  and the global minimum of  $g_\varepsilon^m$  is  $x_0 + \varepsilon u$ . We have:

$$\varepsilon^{-1}s(Tg_\varepsilon^m, Tg) \equiv 1.$$



Next, we will show that  $\theta_\varepsilon^m$  goes to zero fast enough so that  $g_\varepsilon^m$  is very close to  $D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u)$ . We have:

$$\begin{aligned} 0 &= \int (h \circ g_\varepsilon^m - h \circ g) dx \\ &= - \int \left( h \circ g - h \circ ((1 - \theta_\varepsilon^m) D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) + \theta_\varepsilon^m g) \right) dx \\ &\quad + \int (h \circ (\theta_\varepsilon^m D_\delta(g; x_1, y_1) + (1 - \theta_\varepsilon^m)g) - h \circ g) dx, \end{aligned}$$

where both integrals have the same sign. For the first integral by Lemma 3.26 we have:

$$\begin{aligned} &\int \left| h \circ g - h \circ ((1 - \theta_\varepsilon^m) D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) + \theta_\varepsilon^m g) \right| dx, \\ &\leq \int \left| h \circ g - h \circ D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) \right| dx \sim \int \left| D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) - g \right| dx \\ &\leq \xi(\varepsilon) \mu[B_G(x_0 + \varepsilon u; \sqrt{2\xi(\varepsilon)})] = O(\xi(\varepsilon)^{1+d/2}) \end{aligned}$$

The second integral is monotone in  $\theta_\varepsilon^m$  and by Lemma 3.27 we have:

$$\int (h \circ (\theta_\varepsilon^m D_\delta(g; x_1, y_1) + (1 - \theta_\varepsilon^m)g) - h \circ g) dx \sim \theta_\varepsilon^m,$$

thus we have  $\theta_\varepsilon^+ = O(\varepsilon^{\gamma(1+d/2)})$ .

For Hellinger distance we have:

$$\begin{aligned} H(h \circ g_\varepsilon^m, h \circ g) &= H(h \circ ((1 - \theta_\varepsilon^m) D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) + \theta_\varepsilon^m g), h \circ g) \\ &\quad + H(h \circ (\theta_\varepsilon^m D_\delta(g; x_1, y_1) + (1 - \theta_\varepsilon^m)g), h \circ g). \end{aligned}$$

For the first part we can apply Lemma 3.26:

$$\begin{aligned} H^2(h \circ ((1 - \theta_\varepsilon^m) D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) + \theta_\varepsilon^m g), h \circ g) &\leq H^2(h \circ D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u), h \circ g) \\ \lim_{\varepsilon \rightarrow 0} \frac{H^2(h \circ D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u), h \circ g)}{\int (D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) - g)^2 dx} &= \frac{h' \circ g(x_0)^2}{4h \circ g(x_0)} \quad \text{and} \end{aligned}$$

$$\begin{aligned} \int (D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) - g)^2 dx &\leq \xi(\varepsilon)^2 \mu[B_G(x_0; \sqrt{2\xi(\varepsilon)})] \\ &= \xi(\varepsilon)^{2+d/2} \frac{2^{d/2} \mu[S(0, 1)]}{\sqrt{\det G}} \end{aligned}$$

which gives:

$$\begin{aligned} & \limsup_{\varepsilon \rightarrow 0} \varepsilon^{-\gamma(1+d/4)} H(h \circ ((1 - \theta_\varepsilon^m) D_{\xi(\varepsilon)}^*(g; x_0 + \varepsilon u) + \theta_\varepsilon^m g), h \circ g) \\ & \leq C(d) L^{1+d/4} \left[ \frac{h' \circ g(x_0)^4}{h \circ g(x_0)^2 \det G} \right]^{1/4} \end{aligned}$$

where  $S(0, 1)$  is  $d$ -dimensional sphere of radius 1.

For the second part by Lemma 3.27 we obtain:

$$\begin{aligned} & \limsup_{\varepsilon \rightarrow 0} (\theta_\varepsilon^+)^{-2} H^2(h \circ ((1 - \theta_\varepsilon^+)g + \theta_\varepsilon^+ D_\delta(g; x_1, y_1)), h \circ g) < \infty \\ & H(h \circ ((1 - \theta_\varepsilon^+)g + \theta_\varepsilon^+ D_\delta(g; x_1, y_1)), h \circ g) = O(\varepsilon^{\gamma(1+d/2)}). \end{aligned}$$

Thus:

$$\limsup_{\varepsilon \rightarrow 0} \varepsilon^{-\gamma(1+d/4)} H(h \circ g_\varepsilon^+, h \circ g) \leq C(d) L^{1+d/4} \left[ \frac{h' \circ g(x_0)^4}{h \circ g(x_0)^2 \det G} \right]^{1/4}.$$

Finally, we apply Corollary 2.22:

$$\liminf_{n \rightarrow \infty} n^{\frac{2}{\gamma(d+4)}} R_1(n; T, \{p, p_n\}) \geq C(d) L^{-\frac{1}{\gamma}} \left[ \frac{h \circ g(x_0)^2 \det G}{h' \circ g(x_0)^4} \right]^{\frac{1}{\gamma(d+4)}}.$$

Taking the supremum over all  $G \in \mathcal{SC}(g; x_0)$  we obtain the statement of the theorem.  $\square$

### 3.7. Indications of Proofs for Conjectured Rates.

**4. Appendix: some results from convex analysis.** We will use the following general properties of convex sets and convex functions. We use Rockafellar [1970] as a reference.

LEMMA 4.1. *For any convex set  $A$  in  $\mathbb{R}^d$  we have:*

1. *The boundary of  $A$  has Lebesgue measure zero.*
2.  *$A$  has Lebesgue measure zero if and only if it belongs to a  $d-1$  dimensional affine subspace.*
3.  *$A$  has Lebesgue measure  $+\infty$  if and only if it is unbounded and has dimension  $d$ .*

PROOF.

1. If  $A$  is such that  $\text{cl}(A)$  has finite Lebesgue measure then:

$$\begin{aligned}\partial A &\subseteq (1 + \varepsilon) \text{cl}(A) \setminus (1 - \varepsilon) \text{cl}(A), \quad \varepsilon \in (0, 1) \\ \mu[\partial A] &\leq 2\varepsilon\mu[\text{cl}(A)]\end{aligned}$$

and thus  $\mu[\partial A] = 0$ . Since  $\mathbb{R}^d$  is a countable union of closed convex cubes  $B_i$  the result for an arbitrary convex set  $A$  follows from:

$$\partial A \subseteq \bigcup_i \partial(A \cap B_i).$$

2. If  $A$  has dimension  $k \leq d$  then its affine hull  $V$  has dimension  $k$  and  $A$  contains a  $k$ -dimensional simplex  $D$  (Theorem 2.4 Rockafellar [1970]). Then if  $k = d$  we have  $\mu[D] > 0$  and if  $k < d$  we have  $\mu[V] = 0$ .
3. Part 1 implies that it is enough to consider closed convex sets. Part 2 implies that it is enough to prove that an unbounded closed convex set of dimension  $d$  has Lebesgue measure  $+\infty$ . Let  $A$  be such a set; i.e. an unbounded closed convex set. Then  $A$  contains  $d$ -dimensional simplex  $D$  (Theorem 2.4 Rockafellar [1970]) which has non-zero Lebesgue measure. Since  $A$  is unbounded then its recession cone is non-empty (Theorem 8.4 Rockafellar [1970]) and therefore we can choose a direction  $v$  such that  $D + \lambda v \subset A$  for all  $\lambda \geq 0$  which implies  $\mu[A] = +\infty$ .

□

The following lemma shows that convergence of convex sets in measure implies pointwise convergence.

LEMMA 4.2. *Let  $A$  be a convex set in  $\mathbb{R}^d$  such that  $\dim(A) = d$  and  $\text{ri}(A) \neq \emptyset$ . Then:*

1. *Suppose a sequence of convex sets  $B_n$  is such that  $A \subseteq B_n$  and  $\lim \mu[B_n \setminus A] = 0$  then  $\limsup \text{cl}(B_n) = \text{cl}(A)$ ;*
2. *Suppose a sequence of convex sets  $B_n$  is such that  $C_n \subseteq A$  and  $\lim \mu[A \setminus C_n] = 0$  then  $\liminf \text{ri}(C_n) = \text{ri}(A)$ .*

PROOF. By Lemma 4.1 we can assume that  $A$ ,  $B_n$  and  $C_n$  are closed convex sets.

1. If on the contrary, there exists subsequence  $\{k\}$  such that for some  $x \in A^c$  we have  $x \in \cap_{k \geq 1} B_k$  then for  $xA = \text{conv}(\{x\} \cup A)$  we have:

$$\begin{aligned}xA &\subseteq B_k \\ \mu[B_k \setminus A] &\geq \mu[xA \setminus A].\end{aligned}$$

Since  $A$  is closed there exists a ball  $B(x)$  such that  $B(x) \cap A = \emptyset$ . Since  $\text{ri}(A) \neq \emptyset$  there exists a ball  $B(x_0)$  such that  $B(x_0) \subseteq A$  for some  $x_0 \in \text{ri}(A)$ . Then for  $xB = \text{conv}(\{x\} \cup B(x_0))$  we have:

$$\begin{aligned} xB &\subseteq xA \\ \mu[xA \setminus A] &\geq \mu[xB \cap B(x)] > 0. \end{aligned}$$

This contradiction implies  $\limsup B_i = A$ .

2. If on the contrary, there exists a point  $x \in \text{ri}(A)$  and subsequence  $\{k\}$  such that  $x \notin C_k$  for all  $k$  then for each  $C_k$  there exists a half-space  $L_k$  such that  $x \in L_k$  and  $C_k \subseteq L_k^c$ . Let  $B(x)$  be a ball such that  $B(x) \subseteq A$ . We have:

$$\mu[A \setminus C_i] \geq \mu[A \cap L_k] \geq \mu[B(x) \cap L_k] = \mu[B(x)]/2 > 0.$$

This contradiction implies  $\text{ri}(A) \subseteq \liminf C_i$ .

□

Our next lemma shows that the Lebesgue measure of sublevel sets of a convex function grows at most polynomially.

LEMMA 4.3. *Let  $g$  be a convex function and values  $y_1 < y_2 < y_3$  are such that  $\text{lev}_{y_1} g \neq \emptyset$ . Then we have:*

$$(4.24) \quad \mu[\text{lev}_{y_3} g] \leq \left[ \frac{y_3 - y_1}{y_2 - y_1} \right]^d \mu[\text{lev}_{y_2} g].$$

PROOF. By assumption we have:

$$\mu[\text{lev}_{y_3} g] \geq \mu[\text{lev}_{y_2} g] \geq \mu[\text{lev}_{y_1} g] > 0.$$

Let us consider the set  $L$  defined as:

$$L = \{x_1 + k(x - x_1) \mid x \in \text{lev}_{y_2} g\},$$

where  $x_1$  is any fixed point such that  $g(x_1) = y_1$  and

$$k = \frac{y_3 - y_1}{y_2 - y_1} > 1.$$

Then:

$$\mu[L] = k^d \mu[\text{lev}_{y_2} g].$$

and therefore it is enough to prove that  $\text{lev}_{y_3} g \subseteq L$ .

If  $x_3 \in \text{lev}_{y_3} g$  then for  $x_2 = x_1 + (x_3 - x_1)/k$  we have:

$$\begin{aligned} x_3 &= x_1 + k(x_2 - x_1), \\ g(x_2) &\leq (1 - 1/k)g(x_1) + (1/k)g(x_3) = y_2 \end{aligned}$$

and thus  $x_2 \in \text{lev}_{y_2} g$ . □

**COROLLARY 4.4.** *If  $g$  is a convex function then function  $\mu[\text{lev}_y g]$  is continuous on  $(\inf g, \sup g)$ .*

**4.1. Maximal convex minorant.** In this section we describe the convex function  $f_c$  which is in some sense the closest to a given function  $f$ .

**DEFINITION 4.5.** *The maximal convex minorant  $f_c$  of a proper function  $f$  is a supremum of all linear functions  $l$  such that  $l \leq f$ .*

It is possible that  $f$  does not majorate any linear function and then  $f = -\infty$ . However if it is not the case the following properties of the maximal convex minorant hold:

**LEMMA 4.6.** *Let  $f$  be a function and  $f \neq -\infty$  be its maximal convex minorant. Then:*

1.  $f_c$  is a closed proper convex function;
2. if  $f$  is proper convex function then  $f_c$  is its closure;
3.  $f_c \leq f$ ;
4.  $(f_c)^*(y) = \sup_{x \in \mathbb{R}^d} (\langle y, x \rangle - f(x))$ .

**PROOF.** This follows from Corollary 12.1.1 [Rockafellar \[1970\]](#). □

The maximal convex minorant allows us to see an important duality between operations of pointwise minimum and pointwise maximum.

**LEMMA 4.7.** *Let  $f_i$  be a proper convex functions and let  $g = \inf_i f_i$  be the pointwise infimum of  $f_i$ . Then  $(g_c)^* = \sup_i f_i^*$ .*

**PROOF.** This follows from Theorem 16.5 [Rockafellar \[1970\]](#). □

#### 4.2. Subdifferential.

DEFINITION 4.8. *The subdifferential  $\partial h(x)$  of a convex function  $h$  at the point  $x$  is the set of all vectors  $v$  which satisfy the inequality*

$$h(z) \geq \langle v, z - x \rangle + h(x) \quad \text{for all } z.$$

Obviously  $\partial h(x)$  is a closed convex set. It might be empty, but if it is not, the function  $h$  is called *subdifferentiable* at  $x$ .

LEMMA 4.9. *Let  $h$  be a proper convex function then for  $x \in \text{ri dom } h$  subdifferential  $\partial h(x)$  is not empty.*

PROOF. This follows from Theorem 23.4 [Rockafellar \[1970\]](#). □

LEMMA 4.10. *Let  $h$  be a closed proper convex function and  $x$  any point. Then the following conditions on  $x^*$  are equivalent:*

1.  $y \in \partial h(x)$ ;
2.  $l(z) = \langle x^*, z \rangle - h^*(x^*)$  is a support plane for  $\text{epi}(h)$  at  $x$ ;
3.  $h(x) + h^*(x^*) = \langle x^*, x \rangle$ ;
4.  $x \in \partial h^*(y)$ ;
5.  $l(z) = \langle x, z \rangle - h(x)$  is a support plane for  $\text{epi}(h^*)$  at  $x^*$ ;

PROOF. This follows from Theorem 23.5 [Rockafellar \[1970\]](#). □

LEMMA 4.11. *Let  $h_1$  and  $h_2$  be proper convex functions such that  $\text{ri dom } h_1 \cap \text{ri dom } h_2 \neq \emptyset$ . Then  $\partial(h_1 + h_2) = \partial h_1 + \partial h_2$  for all  $x$ .*

PROOF. This follows from Theorem 23.8 [Rockafellar \[1970\]](#). □

#### 4.3. Polyhedral functions.

DEFINITION 4.12. *A polyhedral convex set is a set which can be expressed as an intersection of finitely many half-spaces. A polyhedral convex function is a convex function whose epigraph is polyhedral.*

From Theorem 19.1 [Rockafellar \[1970\]](#) we have that the epigraph of the polyhedral function  $h : \mathbb{R}^d \rightarrow \mathbb{R}$  has finite number of extremal points and faces. We call projections of extremal points the *knots* of  $h$  and projections of the nonvertical  $d$ -dimensional faces the *facets* of  $h$ . Thus the set of knots and the set of facets of polyhedral function are always finite. Moreover, by

Theorem 18.3 [Rockafellar \[1970\]](#) the knots are the extremal points of the facets. Finally, let  $\{C_i\}$  be the set of facets of a polyhedral function  $h$  then:

$$\begin{aligned} \text{dom } h &= \bigcup_i C_i \\ \text{ri}(C_i) \cap \text{ri}(C_j) &= \emptyset, \end{aligned}$$

and on  $\text{dom } h$  we have  $h = \max(l_i)$  where  $l_i$  are linear functions. For each  $C_i$  there exists  $l_i$  such that:

$$C_i = \{x \mid h(x) = l_i(x)\}.$$

LEMMA 4.13. *Let  $f$  be a polyhedral convex function and  $x \in \text{dom } h$  then  $\partial h(x) \neq \emptyset$ .*

PROOF. This follows from Theorem 23.10 [Rockafellar \[1970\]](#). □

LEMMA 4.14. *For the set of points  $x = \{x_i\}_{i=1}^n$  such that  $x_i \in \mathbb{R}^d$  and any point  $p \in \mathbb{R}^n$  consider a family of all convex functions  $h$  with  $\text{ev}_x h = p$ . The unique maximal element  $U_x^p$  in this family is a polyhedral convex function with domain  $\text{dom } U_x^p = \text{conv}(x)$  and the set of knots  $K \subseteq x$ .*

PROOF. Points  $(x_i, p_i)$  and direction  $(0, 1)$  belong to the epigraph of any convex function  $h$  in our family and so does convex hull  $U$  of these points and direction. By construction  $U$  is an epigraph of some closed proper convex function  $U_x^p$  such that  $\text{dom } U_x^p = \text{conv}(x)$ , by Theorem 19.1 [Rockafellar \[1970\]](#) this function is polyhedral, by Corollary 18.3.1 [Rockafellar \[1970\]](#) the set of its knots  $K$  belongs to  $x$  and since  $\text{epi}(U_x^p) = U \subseteq \text{epi}(h)$  we have  $h \leq U_x^p$ . On the other hand, since  $(x_i, p_i) \in U$  we have

$$p_i = h(x_i) \leq U_x^p(x_i) \leq p_i$$

and therefore  $U_x^p(x_i) = p_i$  which proves the lemma. □

LEMMA 4.15. *For the set of points  $x = \{x_i\}_{i=1}^n$ , convex set  $C$  such that  $x_i \in \text{ri}(C)$  and any point  $p \in \mathbb{R}^n$  consider a family of all convex functions  $h$  with  $\text{ev}_x h = p$  and  $C \subseteq \text{dom } h$ . Any minimal element  $L_x^p$  in this family is a polyhedral convex function with  $\text{dom } L_x^p = \mathbb{R}^d$ . For each facet  $C$  of  $L_x^p$ ,  $\text{ri}(C)$  contains at least one element of  $x$ .*

PROOF. For any function  $h$  in our family let us consider the set of linear functions  $l_i$  such that  $l_i(x_i) = h(x_i) = p_i$  and  $l_i \leq h$  and which correspond

to arbitrarily chosen nonvertical support planes for  $\text{epi}(h)$  at  $x_i$ . Then  $L = \max(l_i)$  is polyhedral and since  $l_j(x_i) \leq h(x_i) = p_i$  we have  $L(x_i) = p_i$ . We also have  $\text{dom } L = \mathbb{R}^d$ . If the interior of any facet  $C_i$  of  $L$  does not contain elements of  $x$  we can exclude corresponding linear function  $l_i$  from maximum. For the new polyhedral function  $L' = \max_{j \neq i} l_j$  we still have  $\text{ev}_x L' = p$ . Now, we repeat this procedure until interior of each facet contains at least one element of  $x$  and denote the function we obtained by  $L_x^p$ . If a closed proper convex function  $h$  is such that  $\text{ev}_x h = p$  and  $h \leq L_x^p$ , then consider for any facet  $C_i$  and corresponding linear function  $l_i$  we have  $h \leq l_i$  on  $C_i$  and the supremum of  $h$  on the convex set  $C_i$  is obtained in interior point  $x_j \in x$ . By Theorem 32.1 [Rockafellar \[1970\]](#)  $h \equiv L_x^p$  on  $C_i$ . Thus  $h \equiv L_x^p$  and  $L_x^p$  is the minimal element of our family.  $\square$

LEMMA 4.16. *For linear function  $l(x) = a^T x + b$  the polyhedral set  $A = \{l \geq c\} \cap \mathbb{R}^d$  is bounded if and only if all coordinates of  $a$  are negative. In this case, if  $b \geq c$  the set  $A$  is simplex with vertices  $p_i = ((c - b)/a_i)e_i$  and 0, where  $e_i$  are basis vectors. Otherwise,  $A$  is empty.*

PROOF. If coordinate  $a_i$  is nonnegative then the direction  $\{\lambda e_i\}$ ,  $\lambda > 0$  belongs to the recession cone of  $A$  and thus it is unbounded. If all coordinates  $a_i$  are negative and  $b \leq c$  the set  $A$  is either empty or consists of zero vector 0. Finally, if  $a_i$  are negative and  $b > c$  then for  $x \in A$  we can define  $\theta_i = a_i x_i / (c - b) > 0$ . Then  $1 \geq \sum_i \theta_i$  and  $x = \sum_i \theta_i p_i$ , which proves that  $A$  is simplex.  $\square$

4.4. *Strong convexity.* Following Rockafellar and Wets [Rockafellar and Wets \[1998\]](#) page 565 we say that a proper convex function  $h : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$  is *strongly convex* if there exists a constant  $\sigma$  such that:

$$(4.25) \quad h(\theta x + (1 - \theta)y) \leq \theta h(x) + (1 - \theta)h(y) - \frac{1}{2}\sigma\theta(1 - \theta)\|x - y\|^2$$

for all  $x, y$  and  $\theta \in (0, 1)$ . There is a simple characterization of strong convexity:

LEMMA 4.17. *A proper convex function  $f : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$  is strongly convex if and only if the function  $f(x) - \frac{1}{2}\sigma\|x\|^2$  is convex.*

Since we need a more precise control over the curvature of a convex function we define a generalization of strong convexity based on the characterization above:



DEFINITION 4.18. *We say that a proper convex function  $h : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$  is  $G$ -strongly convex if there exist a point  $x_0$ , a positive semidefinite  $d \times d$  matrix  $G$  and a convex function  $q$  such that:*

$$(4.26) \quad h(x) = \frac{1}{2}(x - x_0)^T G(x - x_0) + q(x) \quad \text{for all } x.$$

Obviously, strong convexity is equivalent to  $\sigma I$ -strong convexity. Note that the definition does not depend on the choice of  $x_0$ .

DEFINITION 4.19. *We say that a proper convex function  $h : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$  is locally  $G$ -strongly convex at a point  $x_0$  if there exist an open neighborhood of  $x_0$ , a positive semidefinite  $d \times d$  matrix  $G$  and a convex function  $q$  such that (4.26) holds for any  $x$  in this neighborhood.*

We can relate  $G$ -strong convexity to the Hessian of a smooth convex function:

LEMMA 4.20. *If a proper convex function  $h : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$  is continuously twice differentiable at  $x_0$  then  $h$  is locally  $(1 - \varepsilon)\nabla^2 h$ -strongly convex for any  $\varepsilon \in (0, 1)$ .*

The last result suggests the following definition:

DEFINITION 4.21. *For a proper convex function  $h : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$  we define a curvature  $\text{curv}_{x_0} h$  at a point  $x_0$  as:*

$$(4.27) \quad \text{curv}_{x_0} h = \sup_{G \in \mathcal{SC}(h; x_0)} \det(G)$$

where  $\mathcal{SC}(h; x_0)$  is the set of all positive semidefinite matrices  $G$  such that  $h$  is locally  $G$ -strong convex at  $x_0$ .

Lemma 4.20 implies that:

LEMMA 4.22. *If a proper convex function  $h : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$  is continuously twice differentiable at  $x_0$  and Hessian  $\nabla^2 h(x_0)$  is positive definite then*

$$(4.28) \quad \text{curv}_{x_0} h = \det(\nabla^2 h(x_0)).$$

## APPENDIX A: NOTATION

|                           |                                                                                                         |
|---------------------------|---------------------------------------------------------------------------------------------------------|
| $\mathbb{R}$              | $= (-\infty, +\infty)$                                                                                  |
| $\overline{\mathbb{R}}$   | $= [-\infty, +\infty]$                                                                                  |
| $\mathbb{R}_+$            | $= [0, +\infty)$                                                                                        |
| $\overline{\mathbb{R}}_+$ | $= [0, +\infty]$                                                                                        |
| $ x $                     | $= \prod_{k=1}^d x_k, \quad x \in \mathbb{R}_+^d$                                                       |
| $\mathcal{C}$             | $= \{f : \mathbb{R}^d \rightarrow \overline{\mathbb{R}} \mid f \text{ closed proper convex function}\}$ |
| $\mathcal{D}$             | $= \{p : \mathbb{R}^d \rightarrow \overline{\mathbb{R}} \mid p \text{ density}\}$                       |
| $\mathcal{G}(h)$          | $= \{h :  g \in \mathcal{C}, h \circ g \in \mathcal{D}\}$                                               |
| $\mathbb{L}_n g$          | $= \mathbb{P}_n h \circ g$                                                                              |
| $\text{ev}_x f$           | $= (f(x_1), \dots, f(x_n)), \quad x_i \in \mathbb{R}^d$                                                 |
| $\text{supp}(f)$          | $= \{x \mid f(x) \neq 0\}$                                                                              |
| $\delta(\cdot \mid C)$    | $= \infty \cdot 1_{C^c} + 0 \cdot 1_C$                                                                  |
| $\text{lev}_y g$          | $= \{x \mid g(x) \leq y\}$                                                                              |
| $\mu[S]$                  | $= \text{Lebesgue measure of } S$                                                                       |
| $\{f \leq a\}$            | $= \{x \in X \mid f(x) \leq a\}$                                                                        |
| $B(x_0; r)$               | $= \{x : \ x - x_0\  < r\}$                                                                             |
| $B_H(x_0; r)$             | $= \{x : (x - x_0)^T H(x - x_0) < r^2\}$                                                                |
| $\text{curv}_x h$         | $= \text{curvature of a convex function } h \text{ at a point } x$                                      |

## ACKNOWLEDGMENTS

This research is a part of the Ph.D. dissertation of the first author at the University of Washington.

## REFERENCES

- AN, M. Y. (1998). Logconcavity versus logconvexity: a complete characterization. *J. Econom. Theory* **80** 350–369.
- AVRIEL, M. (1972).  $r$ -convex functions. *Math. Programming* **2** 309–323.
- BALABDAOUI, F., RUFIBACH, K. and WELLNER, J. A. (2009). Limit distribution theory for maximum likelihood estimation of a log-concave density. *Ann. Statist.* **37** 1299–1331.
- BARAUD, Y. and BIRGÉ, L. (2009). Model selection for functions on high-dimensional spaces. Tech. rep., Université Paris VI. In *Mathematisches Forschungs Institute Oberwolfach Report No. 39/2009, Challenges in Statistical Theory: Complex Data Structures and Algorithmic Optimization*.
- BOBKOV, S. G. (1999). Isoperimetric and analytic inequalities for log-concave probability measures. *Ann. Probab.* **27** 1903–1921.
- BOBKOV, S. G. (2007a). Large deviations and isoperimetry over convex probability measures with heavy tails. *Electron. J. Probab.* **12** 1072–1100 (electronic).
- BOBKOV, S. G. (2007b). On isoperimetric constants for log-concave probability distributions. In *Geometric aspects of functional analysis*, vol. 1910 of *Lecture Notes in Math.* Springer, Berlin, 81–88.
- BOBKOV, S. G. and LEDOUX, M. (2009). Weighted Poincaré-type inequalities for Cauchy and other convex measures. *Ann. Probab.* **37** 403–427.
- BORELL, C. (1975). Convex set functions in  $d$ -space. *Period. Math. Hungar.* **6** 111–136.
- BRASCAMP, H. J. and LIEB, E. H. (1976). On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *J. Functional Analysis* **22** 366–389.
- CORDERO-ERAUSQUIN, D., MCCANN, R. J. and SCHMUCKENSCHLÄGER, M. (2001). A Riemannian interpolation inequality à la Borell, Brascamp and Lieb. *Invent. Math.* **146** 219–257.
- CULE, M. and SAMWORTH, R. (2009). Theoretical properties of the log-concave maximum likelihood estimator of a multidimensional density. Tech. rep., Cambridge University. Available at arXiv:0908.4400v1.
- CULE, M., SAMWORTH, R. and STEWART, M. (2007). Maximum likelihood estimation of a multidimensional log-concave density. Available at arXiv:0804.3989v1.
- DHARMADHIKARI, S. and JOAG-DEV, K. (1988). *Unimodality, convexity, and applications*. Probability and Mathematical Statistics, Academic Press Inc., Boston, MA.
- DONOHU, D. L. and LIU, R. C. (1991). Geometrizing rates of convergence. II, III. *Ann. Statist.* **19** 633–667, 668–701.
- DUDLEY, R. M. (1999). *Uniform central limit theorems*, vol. 63 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge.
- DUENBGEN, L. and RUFIBACH, K. (2009). Maximum likelihood estimation of a log-concave density and its distribution function: basic properties and uniform consistency. Available at arXiv:0709.0334.
- DÜMBGEN, L., HÜSLER, A. and RUFIBACH, K. (2007). Active set and EM algorithms for log-concave densities based on complete and censored data. Tech. rep., University of Bern. Available at arXiv:0707.4643.
- DÜMBGEN, L. and RUFIBACH, K. (2007). Maximum likelihood estimation of a log-concave density and its density and its distribution function. Tech. rep., University of Bern. Available at arXiv:0709.0334.
- FOUGÈRES, P. (2005). Spectral gap for log-concave probability measures on the real line. In *Séminaire de Probabilités XXXVIII*, vol. 1857 of *Lecture Notes in Math.* Springer, Berlin, 95–123.

- GROENEBOOM, P., JONGBLOED, G. and WELLNER, J. A. (2001). Estimation of a convex function: characterizations and asymptotic theory. *Ann. Statist.* **29** 1653–1698.
- IBRAGIMOV, I. A. (1956). On the composition of unimodal distributions. *Teor. Veroyatnost. i Primenen.* **1** 283–288.
- JOHNSON, N. L. and KOTZ, S. (1972). *Distributions in Statistics: Continuous Multivariate Distributions*. Wiley, New York.
- JONGBLOED, G. (2000). Minimax lower bounds and moduli of continuity. *Statist. Probab. Lett.* **50** 279–284.
- KOENKER, R. and MIZERA, I. (2008). Quasi-concave density estimation. *unpublished*.
- MILMAN, E. and SODIN, S. (2008). An isoperimetric inequality for uniformly log-concave measures and uniformly convex bodies. *J. Funct. Anal.* **254** 1235–1268.
- OKAMOTO, M. (1973). Distinctness of the eigenvalues of a quadratic form in a multivariate sample. *Ann. Statist.* **1** 763–765.
- PAL, J. K., WOODROOFE, M. B. and MEYER, M. C. (2007). Estimating a polya frequency function. In *Complex Datasets and Inverse Problems: Tomography, Networks, and Beyond*, vol. 54 of *IMS Lecture Notes-Monograph Series*. IMS, 239–249.
- PRÉKOPA, A. (1973). On logarithmic concave measures and functions. *Acta Sci. Math. (Szeged)* **34** 335–343.
- RINOTT, Y. (1976). On convexity of measures. *Ann. Probability* **4** 1020–1026.
- ROCKAFELLAR, R. T. (1970). *Convex analysis*. Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J.
- ROCKAFELLAR, R. T. and WETS, R. J.-B. (1998). *Variational analysis*, vol. 317 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin.
- RUFIBACH, K. (2006). *Log-concave density estimation and bump hunting for I.I.D. observations*. Ph.D. thesis, Universities of Bern and Göttingen.
- RUFIBACH, K. (2007). Computing maximum likelihood estimators of a log-concave density function. *J. Statist. Comp. Sim.* **77** 561–574.
- SCHUHMACHER, D., HÜSLER, A. and DUEMBGEN, L. (2009). Multivariate log-concave distributions as a nearly parametric model. Tech. rep., University of Bern. Available at arXiv:0907.0250v1.
- UHRIN, B. (1984). Some remarks about the convolution of unimodal functions. *Ann. Probab.* **12** 640–645.
- VAN DER VAART, A. W. and WELLNER, J. A. (1996). *Weak convergence and empirical processes*. Springer Series in Statistics, Springer-Verlag, New York. With applications to statistics.
- WALTHER, G. (2010). Inference and modeling with log-concave distributions. *Statistical Science* **25**.

DEPARTMENT OF STATISTICS, BOX 354322  
 UNIVERSITY OF WASHINGTON  
 SEATTLE, WA 98195-4322  
 E-MAIL: [arseni@stat.washington.edu](mailto:arseni@stat.washington.edu)

DEPARTMENT OF STATISTICS, BOX 354322  
 UNIVERSITY OF WASHINGTON  
 SEATTLE, WA 98195-4322  
 E-MAIL: [jaw@stat.washington.edu](mailto:jaw@stat.washington.edu)